

**DRAFT FOR  
PUBLIC COMMENT**

*Please submit comments to [eli@riverbendsci.com](mailto:eli@riverbendsci.com) or call (707) 832-4206  
Comments due September 18, 2016*

**Upper Klamath Basin  
Nonpoint Source Pollution  
Assessment and Management  
Program Plan**

**Klamath Tribal Water Quality Consortium**

August 2016



**YUROK TRIBE**



**HOOPA VALLEY  
TRIBE**



**KARUK TRIBE**



**RESIGHINI  
RANCHERIA**



**QUARTZ VALLEY INDIAN  
RESERVATION**

# ***DRAFT FOR PUBLIC COMMENT***

*Please submit comments to [eli@riverbendsci.com](mailto:eli@riverbendsci.com) or call (707) 832-4206  
Comments due September 18, 2016*

## **Upper Klamath Basin Nonpoint Source Pollution Assessment and Management Program Plan**

**Klamath Tribal Water Quality Consortium**

Prepared with assistance from:

**Kier Associates**

and

**Riverbend Sciences**

August 2016

# TABLE OF CONTENTS

<b>Table of Contents .....</b>	<b>ii</b>
<b>List of Figures.....</b>	<b>iv</b>
<b>List of Tables.....</b>	<b>iv</b>
<b>1 Overview.....</b>	<b>5</b>
<b>2 Introduction.....</b>	<b>7</b>
2.1 Project Area .....	7
2.1.1 Incorporation by Reference of Consortium Member Tribes’ NPS Plans .....	7
2.2 Tribes Involved .....	9
2.2.1 The Klamath Tribal Water Quality Consortium .....	9
2.2.2 Yurok Tribe .....	10
2.2.3 Hoopa Valley Tribe .....	11
2.2.4 Quartz Valley Indian Reservation .....	11
2.2.5 Karuk Tribe .....	11
2.2.6 Resighini Rancheria.....	11
2.3 Goals and Objectives .....	11
2.4 Public Participation and Governmental Coordination.....	12
<b>3 Methodology.....</b>	<b>12</b>
3.1 Data Sources .....	12
3.2 Previous Assessments .....	15
<b>4 Land Use Summary .....</b>	<b>15</b>
4.1 Climate.....	15
4.2 Waterbodies and Hydrology .....	16
4.3 Fisheries .....	16
4.4 Sub-basins.....	17
4.4.1 Williamson River.....	17
4.4.2 Sprague River (Including Sycan River).....	17
4.4.3 Wood River and Other Tributaries to Upper Klamath Lake.....	17
4.4.4 Upper Klamath Lake Including Agency Lake.....	17
4.4.5 Link River, Lake Ewuana, Keno Reservoir, and Lower Klamath Lake (Link Dam To Keno Dam).....	18
4.4.6 Hydroelectric Reach (Keno Dam to Iron Gate Dam) .....	19
4.4.7 Lost River (Not Included in NPS Project Area) .....	19
4.5 Dams and Diversions .....	19
4.6 Land Use and Ownership.....	20
4.6.1 National Wildlife Refuges .....	23
4.6.2 Forestry.....	23
4.6.3 Agriculture.....	23
4.6.4 Urban.....	23
<b>5 Water Quality Summary.....</b>	<b>24</b>
5.1 Synthesis of Upper Klamath Basin Water Quality Impairments and Links to NPS Pollution .....	24
5.2 Phosphorus.....	25
5.3 Nitrogen .....	27
5.4 Dissolved Oxygen and pH .....	27
5.5 Temperature .....	28
5.6 Cyanobacterial toxins.....	28
5.7 Sediment .....	29
<b>6 Results.....</b>	<b>29</b>
6.1 Beneficial Uses .....	29
6.2 Water Quality Limited Waters.....	32
6.3 Attribution of Impairments to NPS pollution for Each Sub-basin .....	33
<b>7 Discussion .....</b>	<b>34</b>
<b>8 Selection of Best Management Practices.....</b>	<b>34</b>
8.1 Core Participants .....	34
8.2 Public Participation and Governmental Coordination.....	36
<b>9 Existing NPS Control Programs.....</b>	<b>37</b>

9.1	Federal Agencies.....	37
9.1.1	U.S. Bureau of Reclamation.....	37
9.1.2	U.S. Fish and Wildlife Service.....	37
9.1.3	Natural Resources Conservation Service.....	38
9.1.4	U.S. Bureau of Land Management and U.S. Forest Service.....	38
9.1.5	U.S. Environmental Protection Agency.....	38
9.2	The Klamath Tribes of Oregon.....	38
9.2.1	Water Rights and the Upper Klamath Basin Comprehensive Agreement (UKBCA).....	39
9.2.2	Watershed Restoration Plan.....	39
9.3	State Agencies.....	40
9.3.1	Oregon Department of Environmental Quality and Oregon Department of Agriculture.....	40
9.3.2	California State Water Resources Control Board and North Coast Regional Water Quality Control Board.....	40
9.4	Private Companies.....	40
9.4.1	PacifiCorp.....	40
9.5	Non-Profit Organizations.....	41
9.5.1	Klamath Watershed Partnership.....	41
9.5.2	The Nature Conservancy.....	41
9.5.3	Trout Unlimited/Klamath Basin Rangeland Trust.....	41
9.6	Partnership programs.....	42
9.6.1	Klamath Basin Monitoring Program (KBMP).....	42
9.6.2	Upper Klamath Conservation Action Network (UKCAN).....	42
9.6.3	Klamath Regional Conservation Partnership Program (RCPP).....	42
9.6.4	Klamath Basin Restoration Agreement (KBRA).....	42
<b>10</b>	<b>Conclusions.....</b>	<b>43</b>
<b>11</b>	<b>NPS Management Program Plan.....</b>	<b>43</b>
11.1	Management Program Summary.....	43
11.1.1	Administration.....	44
11.2	Funding Sources.....	45
11.2.1	Federal.....	45
11.2.2	State.....	46
11.2.3	Private.....	47
11.3	Categories of Nonpoint Source Pollution.....	47
11.4	Tasks and BMPs.....	48
11.4.1	Agriculture.....	51
11.4.2	Hydromodification and Habitat Alteration.....	54
11.4.3	Forestry.....	56
11.4.4	Urban.....	57
11.4.5	Other.....	58
<b>12</b>	<b>References.....</b>	<b>59</b>
<b>13</b>	<b>Acronyms and Abbreviations List.....</b>	<b>73</b>
<b>APPENDIX A: Oregon Department of Environmental Quality List of Impaired Waterbodies.....</b>		<b>A1</b>
<b>APPENDIX B: California List of Impaired Waterbodies.....</b>		<b>B1</b>

## LIST OF FIGURES

Figure 1. Map showing location of Klamath Basin. The dotted red line is the outline of the NPS project area .....	6
Figure 2. Map showing location of the Consortium’s Upper Klamath Basin Nonpoint Source (NPS) Assessment and Management Program Plan (AMPP) area boundary .....	8
Figure 3. Map of land ownership in the Klamath Basin. The dotted red line is the outline of the NPS project area. Map adapted from Stillwater Sciences et al. (2012).....	10
Figure 4. Map of monitoring stations in the Klamath River Basin, produced by the Klamath Basin Monitoring Program (KBMP).....	14
Figure 5. Map of historic and current Lower Klamath Lake (LKL) and wetlands. Source: USBR (2005).....	18
Figure 6. Land cover in the Upper Klamath basin. Data are from the Multi-Resolution Land Characteristics Consortium for the year 2011.....	21
Figure 7. Map of land ownership in the Upper Klamath Basin. The dotted red line is the outline of the NPS project area. Map adapted from Stillwater Sciences et al. (2012).....	22
Figure 8. Seasonal nutrient mechanisms in Upper Klamath Lake at a watershed scale. Figure copied from Stillwater Sciences et al. (2013). .....	24
Figure 9. Effect of Upper Klamath Lake algal blooms on Keno Reservoir and the Klamath River downstream. Figure copied from Stillwater Sciences et al. (2013).....	25
Figure 10. Relative contributions of tributaries and other sources to external total phosphorus (TP) load, inflow water volume, and drainage area to Upper Klamath Lake for hydrologic years 1992-2010. Figure adapted from Stillwater Sciences et al. (2012) based on data from Walker et al. (2012).....	27

## LIST OF TABLES

Table 1. Selected agencies and organizations with water quality monitoring programs within the Consortium’s NPS AMPP area and the mainstem Klamath River downstream. Table also includes instructions for accessing data. ....	13
Table 2. Land cover types in Upper Klamath Basin NPS AMPP area. ....	20
Table 3. ODEQ Designated Beneficial Uses for the Upper Klamath River and other Basin waters in Oregon.....	30
Table 4. NCRWQCB Designated Beneficial Uses for the Middle Klamath River from Hornbrook upstream to Copco Lake. ....	31
Table 5. Assessment categories of water body support of designated beneficial uses. ....	31
Table 6. Sub-basin summary of impairments and causes of NPS pollution. For the sake of simplicity and usefulness, the table does not list every impairment for every specific waterbody but rather focuses on the most important impairments and causes at the sub-basin level. ....	33
Table 7. Core participants for BMPs.....	35
Table 8. Management program initiation timeline and annual milestones. ....	45
Table 9. NPS implementation schedule by fiscal year (July 1 – June 30). Tasks are sorted first by NPS category and then by goal. Key to NPS categories: Agriculture (AGR), Hydromodification and Habitat Alteration (HHA), Forestry (FOR), Urban (URB), and Other (OTH).....	49
Table A10. Oregon Department of Environmental Quality (ODEQ) list of waterbodies in the Upper Klamath Basin designated as impaired under section 303(d) of the Clean Water Act.....	A1
Table A11. Oregon Department of Environmental Quality (ODEQ) list of impaired lakes and reservoirs in the Upper Klamath Basin.....	A18
Table B12. California 303(d) list for the portion of the Klamath River from the Oregon border downstream to Scott River, USEPA approved 2012. ....	B1

## 1 OVERVIEW

The Consortium produced this Nonpoint Source (NPS) Assessment and Management Program Plan (AMPP) to address water quality issues in the Upper Klamath Basin which affect the Lower Klamath Basin (Figure 1). Water quality problems in the Upper Klamath Basin and its tributaries have been well documented in the Oregon Department of Environmental Quality Total Maximum Daily Loads (TMDLs) for Upper Klamath Lake (ODEQ 2002) and Upper Klamath and Lost rivers (ODEQ 2010b), California North Coast Regional Water Quality Control Board Klamath River TMDL (NCRWQCB 2010), evaluations of techniques for water quality improvement (Stillwater Sciences et al. 2012, 2013), an Environmental Impact Statement/Report for the proposed removal of the Klamath Hydroelectric Project (US DOI and CDFG 2012), and numerous other studies by federal, tribal, and state agencies. At Iron Gate Dam near the California border, the Klamath River water is often of insufficient quantity and poor quality to meet the needs of fish, wildlife, and humans. To address this problem, the Consortium's goal is to improve land and water management in the Upper Klamath Basin area to improve the quality of water entering the Lower Klamath Basin.

This NPS AMPP covers the portion of the Klamath Basin that is upstream of Iron Gate Dam near Hornbrook, CA, excepting the Lost River and Butte sub-basins. This area was chosen for this assessment because water quality impacts to water quality on the Klamath River primarily occur upstream of this location.

In the NPS assessment (Sections 2 through 10 below), the Consortium reviews available scientific information regarding the causes and potential solutions for the NPS pollution in the Upper Klamath Basin, including a review of existing NPS management efforts. The assessment identified categories of NPS pollution that are likely impacting water quality and then ranked them based on their relative importance. The following two categories are high priority, due to the widespread extent and severity of impacts:

- Agriculture
- Hydromodification and Habitat Alteration

The following two categories are much lower priority given their lesser contribution to NPS pollution within the NPS AMPP area:

- Forestry
- Urban

The NPS management plan (Section 11) then proposes a schedule (Table 9) of tasks and best management practices (BMPs) and identifies potential funding sources (Section 0) to address the causes of the NPS pollution.

The entire NPS AMPP area is outside the reservations of the Consortium member Tribes; however, some of the area is within the ancestral territory of Shasta Indians who are enrolled members of QVIR. Consortium member Tribes have limited legal authority to mandate changes in land and water management; therefore, the Consortium's NPS AMPP relies on voluntary measures and collaboration with entities already doing work in the area. Rather than "re-invent the wheel," the Consortium intends to support organizations and programs that are already implementing effective projects to restore water quality. The Consortium will continue to develop working relationships with entities addressing water quality issues in the Upper Klamath Basin.




Figure 1. Map showing location of Klamath Basin. The dotted red line is the outline of the NPS project area. Map adapted from Stillwater Sciences et al. (2012).

## 2 INTRODUCTION

The Consortium produced this Nonpoint Source (NPS) Assessment and Management Program Plan (AMPP) to address water quality issues in the Upper Klamath Basin which affect the Lower Klamath Basin (Figure 1). The Consortium used the U.S. EPA's (2010) *Handbook for Developing and Managing Tribal Nonpoint Source Pollution Programs Under Section 319 of the Clean Water Act* and the Yurok Tribe's (2014) *NPS Assessment and Management Program Plan* as the primary templates for this NPS AMPP.

### 2.1 PROJECT AREA

The Klamath River Basin (Figure 1) is 12,680 square miles in area and originates in southern Oregon and extends to northern California before it reaches the Pacific Ocean at Requa in Del Norte County, CA. Forty-four percent of the watershed lies within Oregon while the remaining 56 percent lies within California (ODEQ 2010b). This NPS AMPP covers the portion of the Klamath Basin that is upstream of Iron Gate Dam near Hornbrook, CA, excepting the Lost River and Butte sub-basins (Figure 2). This area was chosen because it is the primary source of impacts to Klamath River water quality which extend all the way downstream to the Klamath Estuary.

While typically considered part of the Klamath Basin, the Lost River is excluded from the project area because it does not naturally connect to the Klamath River. Groundwater elevations suggest that groundwater from the Butte sub-basin drains to Lower Klamath Lake and the Klamath River (Gannett et al. 2007), but there is no surface water connection so the Butte sub-basin is not included in the project area.

The tributaries of the Klamath River originate at the southeast end of the Cascade Mountains in Oregon and converge into Upper Klamath Lake from which the Klamath River flows southwest into California and confluences with the Pacific Ocean near Requa, CA. In Oregon, major tributaries include the Williamson, Wood, Sprague, and Sycan rivers. In California, major tributaries include the Shasta, Scott, Salmon, and Trinity rivers. Six dams are present on the Klamath River between Upper Klamath Lake and the Shasta River. This NPS AMPP focuses on the Upper Klamath River from Iron Gate Dam upstream, which includes multiple tributaries to the River (i.e., Wood, Williamson, Sprague, and Sycan rivers), lakes (Upper Klamath, Agency, and Lower Klamath lakes), and reservoirs (Keno, Copco, Iron Gate, and JC Boyle). The Klamath Straits Drain and other major irrigation and ditch systems are also included.

#### 2.1.1 INCORPORATION BY REFERENCE OF CONSORTIUM MEMBER TRIBES' NPS PLANS

Four of the five Consortium members have a U.S. EPA-approved NPS plan for their reservations, ancestral territory, or relevant watersheds (Karuk Tribe 2003, HVTEPA 2008, QVIR 2008, YTEP 2014, HVTEPA no date). The Consortium hereby incorporates those previous NPS plans into this new Consortium NPS AMPP by reference. This incorporation by reference will allow the Consortium to also be eligible to apply for USEPA Clean Water Action Section 319 funding for geographic areas covered by individual Tribes' NPS plans as well as the Upper Klamath Basin.






Figure 2. Map showing location of the Consortium’s Upper Klamath Basin Nonpoint Source (NPS) Assessment and Management Program Plan (AMPP) area boundary.

## 2.2 TRIBES INVOLVED

### 2.2.1 THE KLAMATH TRIBAL WATER QUALITY CONSORTIUM

The Klamath Tribal Water Quality Consortium (Consortium) is comprised of five federally recognized Tribes, including the Yurok Tribe, Hoopa Valley Tribe, Karuk Tribe, Quartz Valley Indian Reservation and Resighini Rancheria. These Tribes reside within the California portion of the Klamath River Basin on reservation, trust, and fee lands (Figure 3). The federally recognized Klamath Tribes of Oregon (see Section 9.2) is not a member of the Consortium but cooperate with the California tribes on water quality-related issues.

In 2002, over 34,000 adult salmon perished in a single event on the lower Klamath River, representing 19% of the adult salmon returning to spawn. The cause of the die-off was fish pathogens that overtook salmon weakened by low, warm water flows in the Klamath River (USFWS 2003a, 2003b). From this occurrence, the California Klamath Tribes saw a need to protect their threatened cultural resources through engagement of environmental management, monitoring, and policy development. Therefore, in 2003, the tribes formed the Klamath Basin Tribal Water Quality Work Group that worked collectively on a variety of water quality issues with success. In 2015, tribal collaborations were formalized by creating the Consortium. The Consortium serves to enhance the ability of tribes to work with state and federal partners to restore the Klamath Basin and its resources Consortium members depend upon (KTWQC 2015).




Figure 3. Map of land ownership in the Klamath Basin. The dotted red line is the outline of the NPS project area. Map adapted from Stillwater Sciences et al. (2012).

### 2.2.2 YUOK TRIBE

The Yurok Tribe is California’s largest tribe, with nearly 5,000 enrolled members<sup>1</sup>. The Yurok Reservation lands extend from one mile on each side from the mouth of the Klamath River upriver for a distance of 46 miles, where the Yurok Reservation shares a border with the Hoopa Tribe. The Yurok Tribe maintains jurisdiction over waters that flow into and through the reservation for water quality purposes, regardless of the geographic origin of the water sources. The Yurok Tribe Environmental Program (YTEP) developed a Water Quality Control Plan for

<sup>1</sup> www.yuroktribe.org accessed January 2016

reservation waters in 2004 and in 2014 developed a Non-Point Source Assessment and Management Program Plan (YTEP 2004, 2014).

### 2.2.3 HOOPA VALLEY TRIBE

The Hoopa Valley Reservation has approximately 3,346 members and is the largest reservation in California, encompassing 93,702 acres. Hoopa tribal lands are located in the northeastern corner of Humboldt County in Northern California. The reservation is bisected in a north-south direction by the Trinity River while the Klamath River flows east-west through a portion of the northeastern part of the reservation (HVTEPA 2008).

The Hoopa Valley Tribe obtained treatment as a state (TAS) status by EPA in 1996 for purposes of water pollution control (<http://www.epa.gov/wqs-tech/epa-approvals-tribal-water-quality-standards>). From this, the Hoopa Valley Tribe maintains jurisdiction for water quality purposes over all waters that flow into and through the reservation, regardless of the geographic origins of water sources. The Hoopa Valley Tribal Environmental Protection Agency (HVTEPA) developed a Water Quality Control Plan for the Hoopa Valley Reservation in 2002 and the document was amended in 2008 (HVTEPA 2008).

### 2.2.4 QUARTZ VALLEY INDIAN RESERVATION

The Quartz Valley Indian Reservation (QVIR) is located in Quartz Valley, which is in the Scott River sub-basin of the Klamath Basin in California. (he Quartz Valley Tribal Environmental Program developed an EPA-approved NPS plan (QVIR 2008) and Quality Assurance Project Plan (QVIR 2016). The original reservation boundary is 604 acres, but much of the land within it is privately owned by people who are not members of QVIR. QVIR has approximately 250 enrolled members, most of which live on the reservation. Some enrolled members of QVIR are Shasta Indians whose ancestral territory includes portions of the NPS AMPP area.

### 2.2.5 KARUK TRIBE

The Karuk Tribe does not possess reservation lands but resides on 1,168 acres of tribal trust and private domain allotments that include properties along the middle portion of the Klamath River and its tributaries in Northern California. The Karuk Tribe Department of Natural Resources developed a Water Quality Control plan in 2002 and updated the document in 2014 (Karuk Tribe 2014).

### 2.2.6 RESIGHINI RANCHERIA

The Resighini Rancheria Reservation is situated on 228 acres located along the south bank of the Klamath River. The Resighini Rancheria Environmental Protection Authority (REPA) developed a Water Quality Ordinance in 2002 (updated in 2006) which sets water quality standards for the reservation (REPA 2006). In 2010, the REPA developed a surface water sampling and analysis plan (REPA 2010).

## 2.3 GOALS AND OBJECTIVES

Although the Upper Klamath Basin does not contain land held by Consortium member Tribes, the Consortium seeks to apply BMPs and watershed improvements to improve water quality which impacts downstream tribal waters, fisheries, and quality of life. It is not the intention of

the Consortium to initiate new unilateral projects, but rather to support established programs with sustained records of success.

The Consortium's objectives are to enhance and improve the qualities of water and aquatic systems within the Klamath River Basin through:

- Assisting in restoration programs already in place;
- Providing funding to support programs that improve the natural functioning of watersheds in the Upper Klamath River; and
- Provide technical support where needed for project development.

## **2.4 PUBLIC PARTICIPATION AND GOVERNMENTAL COORDINATION**

NPS pollution is a community-wide issue and successful implementation of the NPS AMPP will rely upon relationships between the Consortium, our partner entities, and the public, including tribal and non-tribal community members. Therefore, the Consortium sought public input on this NPS AMPP by engaging public agencies that have a role in managing or protecting natural resources. The Consortium did an oral presentation on the NPS AMPP at the spring 2016 Klamath Basin Monitoring Program (KBMP) meeting which was attended by approximately 50 people, primarily representatives of entities involved in Klamath Basin water quality issues. The Consortium and its consultants conducted phone meetings with many partner organizations to get input on the NPS AMPP, and followed up with email correspondence.

In addition, the Consortium made a draft version of this document available for a 30-day public comment period starting August 19, 2016. Public notice was made by announcing the release of the document in each Consortium Tribe's newsletter, official website, and social media sites. The public notice was also listed on the KBMP website. The Consortium will review all comments that were received and consider them thoroughly before making appropriate changes to the document.

## **3 METHODOLOGY**

### **3.1 DATA SOURCES**

Monitoring data and summaries are available for public use on the Klamath Basin Monitoring Program (KBMP, website: [www.kbmp.net](http://www.kbmp.net)). KBMP is working to develop a monitoring program for the Klamath River basin that does not replace individual water quality monitoring efforts, but expands coordinated monitoring to benefit long-term collaboration. KBMP's goal is to include all agencies and organizations that engage in water quality monitoring in the Klamath River basin. Water quality monitoring in the basin is performed by over 20 tribal, federal, state, and county agencies and private and nonprofit groups throughout the entire Klamath Basin (Figure 4). Table 1 provides some links to access water quality data<sup>2</sup>. Individual tribes in the Consortium have their own environmental departments with expertise to derive, analyze,

---

<sup>2</sup> Additional monitoring data links are available at <http://www.kbmp.net/maps-data/links-data-reports>

implement, and evaluate management decisions. The majority of the data collection programs were initiated in 2002 after the fish kill that occurred in the Lower Klamath River.

Table 1. Selected agencies and organizations with water quality monitoring programs within the Consortium’s NPS AMPP area and the mainstem Klamath River downstream. Table also includes instructions for accessing data. Entities that provide only a portion of their data online are listed as “Request.”

Category	Agency/organization	Data Access
Tribal	Klamath Tribes	Request, <a href="http://www.ceden.org">http://www.ceden.org</a>
	Yurok Tribe	Request, <a href="http://www.epa.gov/storet/">http://www.epa.gov/storet/</a> , <a href="http://exchange.yuroktribe.nsn.us/lrgsclient/stations/stations.html">http://exchange.yuroktribe.nsn.us/lrgsclient/stations/stations.html</a>
	Karuk Tribe	Request, <a href="http://www.epa.gov/storet/">http://www.epa.gov/storet/</a> , <a href="http://waterquality.karuk.us">waterquality.karuk.us</a> ,
	Hoopla Tribe	Request, <a href="http://www.epa.gov/storet/">http://www.epa.gov/storet/</a>
	Quartz Valley Indian Reservation	Request, <a href="http://www.epa.gov/storet/">http://www.epa.gov/storet/</a> , <a href="http://www.ceden.org">http://www.ceden.org</a> ,
	Resighini Rancheria	Request, <a href="http://www.epa.gov/storet/">http://www.epa.gov/storet/</a> ,
	Federal	U.S. Bureau of Reclamation
U.S. Bureau of Land Management		Request
U.S. Fish and Wildlife Service		Request
U.S. Forest Service (Fremont-Winema, Klamath, Modoc)		Request, <a href="http://www.fs.fed.us/rm/boise/AWAE/projects/NorWeST/StreamTemperatureDataSummaries.shtml">http://www.fs.fed.us/rm/boise/AWAE/projects/NorWeST/StreamTemperatureDataSummaries.shtml</a>
U.S. Geological Survey		<a href="http://waterdata.usgs.gov/nwis">http://waterdata.usgs.gov/nwis</a> , <a href="http://or.water.usgs.gov/grapher">http://or.water.usgs.gov/grapher</a> , <a href="http://or.water.usgs.gov/projs_dir/klamath_ltmon/">http://or.water.usgs.gov/projs_dir/klamath_ltmon/</a>
State	California Department of Water Resources	<a href="http://cdec.water.ca.gov/">http://cdec.water.ca.gov/</a> , <a href="http://www.water.ca.gov/waterdatalibrary">http://www.water.ca.gov/waterdatalibrary</a>
	North Coast Regional Water Quality Control Board	Request, <a href="http://www.ceden.org">http://www.ceden.org</a>
	Oregon Department of Environmental Quality	<a href="http://deq12.deq.state.or.us/lasar2">http://deq12.deq.state.or.us/lasar2</a>
Private	PacifiCorp	<a href="http://www.pacificorp.com/es/hydro/hl/kr.html">http://www.pacificorp.com/es/hydro/hl/kr.html</a> , <a href="http://www.kbmp.net/collaboration/klamath-hydroelectric-settlement-agreement-monitoring">http://www.kbmp.net/collaboration/klamath-hydroelectric-settlement-agreement-monitoring</a> ,
Non-Profit	Trout Unlimited (Klamath Basin Rangeland Trust)	Request
	The Nature Conservancy	Request




Figure 4. Map of monitoring stations in the Klamath River Basin, produced by the Klamath Basin Monitoring Program (KBMP).

### 3.2 PREVIOUS ASSESSMENTS

The Consortium relies on the monitoring efforts of the above-mentioned agencies and groups to determine the quality of waters and the sources of impairments. The large area of land the Klamath River Basin encompasses makes initiating a new and separate monitoring program inefficient and redundant. The Consortium assumes organizations (Table 4) perform effective quality control and assurances with data collection and analysis.

In addition to the basin-wide monitoring efforts, the Consortium relies heavily on a large number of previous analyses and syntheses of water quality information from reputable sources, including federal, state, tribal, academic, consultant, and non-profit organizations. These documents are cited in the appropriate sections below. A few of the documents most essential to the development of this NPS AMPP are: Total Maximum Daily Loads (TMDLs) for Upper Klamath Lake (ODEQ 2002), Upper Klamath and Lost Rivers (ODEQ 2010b), and Klamath River (NCRWQCB 2010); lists of impaired water bodies from the states of Oregon and California; nutrient budgets for Upper Klamath Lake (Walker et al. 2012) and its tributaries (Walker et al. 2015); studies of Upper Klamath Lake ecology and water quality (Kann 1998, Kann and Smith 1999, Bradbury et al. 2004, Eilers et al. 2004, Kann and Welch 2005); evaluations of techniques for water quality improvement (Stillwater Sciences et al. 2012, 2013); Environmental Impact Statement/Report for the proposed removal of the Klamath Hydroelectric Project (US DOI and CDFG 2012); reports regarding water quality models for Keno Reservoir (Sullivan et al. 2011, 2013, 2014) and Upper Klamath Lake (Wood et al. 2008, Wherry et al. 2015).

## 4 LAND USE SUMMARY

### 4.1 CLIMATE

The following excerpt from Stillwater Sciences et al. (2012) summarizes the Klamath Basin's climate:

“Annual precipitation in the Klamath Basin ranges from 15–150 inches per year, with drier conditions (15–40 inches per year) at the higher elevations of the upper basin (i.e., greater than 1,219 m [4,000 ft]) and wetter conditions (40–150 inches per year) at lower elevations and in coastal areas of the lower basin. The upper basin receives rain and snow during the late fall, winter and spring, with most winter precipitation falling as snow. The Upper Klamath Lake system freezes over intermittently from November through February, and can remain frozen for several months duration. Midwinter rains can occur in the lower-elevation areas (i.e., 914–1,219 m [3,000–4,000 ft]) of the upper basin, but due to a rain shadow effect of the Cascade Mountains annual rainfall is variable throughout this portion of the Klamath Basin (Risley and Laenen 1999), and ranges from a mean annual precipitation (1961–1990) level of 166.1 cm (65.4 in) at Crater Lake National Park in the Cascade Range to 34.3 cm (13.5 in) at Klamath Falls (Gannett et al. 2007).”



## 4.2 WATERBODIES AND HYDROLOGY

The Klamath River's water originates in tributaries that flow from the southeast boundary of the Cascade Mountains and from mountains that form the southwestern edge of the Great Basin (Figure 1, Figure 2). These tributaries converge into the large, shallow Upper Klamath Lake and its northern arm which is known as Agency Lake. The Wood River flows from the flanks of the Cascade Mountains into Agency Lake and then into Upper Klamath Lake. Another major tributary, the Williamson River, originates in the Winema National Forest and flows directly into Upper Klamath Lake. Two major streams contribute flow to the Williamson River, including the Sprague River and its tributary, the Sycan River, both of which originate in the forested mountains of the Fremont National Forest. The Williamson, Sprague, and Sycan rivers contribute 79% of the drainage area for Upper Klamath Lake (ODEQ 2002). Water levels in Upper Klamath Lake are regulated by Link Dam. The outlet of Upper Klamath Lake flows into the Link River, which flows 1.2 miles before entering Lake Ewauna. The Klamath River proper begins at the outlet of Lake Ewauna, which is currently part of Keno Reservoir impounded by Keno Dam.

Below Keno Dam are an additional four dams on the mainstem Klamath River which form the core of the Klamath Hydroelectric Project (KHP) (see Section 4.5 for additional information on dams on the Klamath River). Between Keno Dam and Iron Gate Dam (the lowest of the mainstem dams), the Klamath River flows increase substantially due to contributions from springs below JC Boyle Dam and tributaries including Spencer Creek, Shovel Creek, Fall Creek, and Jenny Creek. The Lost River does not naturally connect to the Klamath River but is connected via the Lost River Diversion Channel which diverts water from the Klamath River during irrigation season (June-September) and contributes irrigation runoff to the Klamath River via a series of pumps and other infrastructure that operate the remainder of the year. The Lost River sub-basin is not part of this assessment.

Due to permeable geology, streams in the Upper Klamath Basin have relatively high summer base flows. Average groundwater discharge into streams of the Upper Klamath Basin upstream from Iron Gate Dam is about 2,400 cfs (1.8 million acre-ft/yr) (Gannett et al. 2007).

## 4.3 FISHERIES

The Consortium Tribes in California conduct robust, active fisheries programs, which help sustain tribal subsistence fishing practices that are culturally and economically significant to the Tribes. Tribal fishing centers on Chinook salmon (*Oncorhynchus tshawytscha*), but some coho salmon (*O. kisutch*) and steelhead (*O. mykiss*) are also caught. The Klamath Tribes in Oregon historically relied on the culturally significant fisheries of Chinook salmon, Lost River sucker (*Deltistes luxatus*) and Shortnose sucker (*Chasmistes brevirostris*).

Historically, salmon have spawned, reared, and migrated to and from the tributaries in the basin. In 1918 with the installment of Copco 1 dam, fisheries habitat and spawning grounds upstream of the dam were entirely cut off from the lower Klamath River (Hamilton et al. 2005). Iron Gate Dam was built downstream in 1962, further reducing the habitat accessible to anadromous salmonids. Presently, over 420 miles of salmonid habitat in the Upper Klamath Basin is inaccessible due to the dams (U.S. DOI and CDFG 2012).

The Upper Klamath River, lakes and reservoirs contain a variety of coldwater and warmwater fisheries. Coldwater fish species found in the basin include native redband trout (*Oncorhynchus mykiss*) and bull trout (*Salvelinus confluentus*) and non-native Eastern brook trout (*Salvelinus*

*fontinalis*) and brown trout (*Salmo trutta*). Warmwater fish species include brown bullhead (*Ameiurus nebulosus*), largemouth bass (*Micropterus salmoides*), pumpkinseed (*Lepomis gibbosus*) and yellow perch (*Perca flavescens*). Federally listed endangered or threatened fish species present in the Upper Klamath Basin include bull trout, Lost River sucker, and shortnose sucker (ODEQ 2002).

#### 4.4 SUB-BASINS

##### 4.4.1 WILLIAMSON RIVER

The Williamson River (4<sup>th</sup> field HUC 18010201) watershed is approximately 1420 mi<sup>2</sup> and has an elevation range of 9,182 feet at the summit of Mount Thielsen to the Williamson River Delta (4,143 feet elevation), which is located on the northeast shores of Upper Klamath Lake. The Williamson River flows in a horseshoe shape, heading north, west, and finally south to Upper Klamath Lake. The Williamson River watershed is relatively flat as over 70% of the watershed has a slope of less than 8% and 50% of the watershed has a slope below 3% (David Evans and Associates 2005). The Klamath Marsh National Wildlife Refuge's 40,000 acres of wet meadows and open water wetlands lies in the middle of the Williamson River watershed and is managed by U.S. Fish and Wildlife Service (USFWS). The Williamson River flows into and out of the marsh. The watershed geology is predominately volcanic in origin, and consists of ash, pumice, and basalt. Due to the porous geology, many tributaries on the west side of the watershed flow subsurface before reaching the Williamson River as springs (David Evans and Associates 2005).

##### 4.4.2 SPRAGUE RIVER (INCLUDING SYCAN RIVER)

The Sprague River (4<sup>th</sup> field HUC 18010202) watershed is 1580 mi<sup>2</sup>, originates on the forested slopes within the Fremont-Winema National Forest at elevations above 7,000 feet and flows south and west towards its confluence with the Sycan River near the town of Beatty, Oregon. From here, the Sprague River flows west to its confluence with the Williamson River. The Sycan River originates at Winter Ridge (6,700 feet), which forms the eastern edge of the Klamath River watershed. The Sycan River flows northwest into Sycan Marsh and then south to its confluence with the Sprague River. Over half of the Sprague River watershed is owned by federal or state agencies. The remaining land is privately owned and managed for commercial timber or rangeland agriculture. The private agricultural lands (pasture and hay) are located in the alluvial valleys along the mainstem Sprague River (O'Connor et al. 2015).

##### 4.4.3 WOOD RIVER AND OTHER TRIBUTARIES TO UPPER KLAMATH LAKE

The Wood River flows from the east side of Crater Lake National Park and terminates at Agency Lake near Chiloquin, Oregon. The Wood River meanders through predominantly agricultural lands consisting of irrigated pasture. Unlike the Williamson and Sprague Rivers, the Wood River watershed does not have its own 4<sup>th</sup> field HUC and is considered by USGS as part of the Upper Klamath Lake hydrologic unit (18010203). Additional tributaries to Upper Klamath Lake include Sevenmile Creek/Canal and Fourmile Creek/canal.

##### 4.4.4 UPPER KLAMATH LAKE INCLUDING AGENCY LAKE

Upper Klamath Lake is a large (235 km<sup>2</sup>) and shallow (mean depth 2 m) hyper-eutrophic lake. The broad, shallow morphology of Upper Klamath Lake affects water quality by enabling solar heating of water temperatures and wind-driven mixing (Wood et al. 2008). The size of Upper

Klamath Lake has been reduced from its historic extent by the agricultural draining of wetlands surrounding the lake (Snyder and Morace 1997). The northern arm of Upper Klamath Lake is known as Agency Lake. Agency Lake is a smaller lake (35 km<sup>2</sup>) and is shallow and eutrophic. Levee breaching in the Williamson River delta in 2007 and 2008 has increased connectivity between Upper Klamath Lake and Agency Lake (Wood et al. 2014).

#### 4.4.5 LINK RIVER, LAKE EWUANA, KENO RESERVOIR, AND LOWER KLAMATH LAKE (LINK DAM TO KENO DAM)

Link River is a 1.2-mile reach of high-gradient river running from Link Dam (the outlet of Upper Klamath Lake) to Lake Ewauna (Figure 2 and Figure 5). The reach of the Klamath River downstream of Lake Ewauna is currently impounded by Keno Dam which forms Keno Reservoir, but was formerly part of Lower Klamath Lake. Lower Klamath Lake has been nearly drained from its original size of approximately 80,000 acres of wetlands and open water (Weddell 2000) (Figure 5). Prior to the construction of the railroad across the Klamath Straits, water from the Klamath River would fill Lower Klamath Lake during spring runoff and then reverse direction and flow back to the river in summer and fall (Weddell 2000). The northern half of the former lakebed is primarily private agricultural lands while the southern half is primarily in the Lower Klamath National Wildlife Refuge, part of which is also used for agricultural purposes. Water is delivered to the wildlife refuge and agricultural lands from the Klamath River via the Ady Canal and North Canal, and from Tule Lake via the tunnel through Sheepy Ridge. Keno Reservoir is held at nearly static elevation to optimize the system of irrigation diversions and agricultural drains.




Figure 5. Map of historic and current Lower Klamath Lake (LKL) and wetlands. Source: USBR (2005).

#### 4.4.6 HYDROELECTRIC REACH (KENO DAM TO IRON GATE DAM)

Downstream of Keno Dam, the Klamath River enters a steep canyon which includes some reaches that are impounded by dams (see Section 4.5). The land use is primarily private timberlands with some U.S. Bureau of Land Management and U.S. Forest Service land and a few small ranches (Figure 7).

#### 4.4.7 LOST RIVER (NOT INCLUDED IN NPS PROJECT AREA)

Historically, the Lost River received water from the Klamath River during high flow events when water would spill through the Lost River Slough (Figure 5) (NRC 2004). The waters would flow toward Tule Lake, the terminus of the Lost River. The Lost River was formerly a closed basin with no natural surface outflow connectivity to other watersheds. The Lost River Diversion channel now connects the Klamath River to the Lost River, and water can flow in either direction, depending on the dual water management goals of drainage and irrigation. Water is also pumped from the D Plant at Tule Lake west to Lower Klamath Lake through a tunnel under Sheepy Ridge. The contribution of the Lost River subwatershed to the overall quality of the Klamath River is seasonally variable. The amount of water pumped through the D Plant has declined significantly in the past decade, potentially due to groundwater depletion within the Tule Lake sub-basin (Pischel and Gannett 2015).

The Lost River is highly altered from its historical condition, as eloquently summarized by USFWS (2001):

“The Lost River can perhaps be best characterized as an irrigation water conveyance, rather than a river. Flows are completely regulated, it has been channelized in one six- mile reach, its riparian habitats and adjacent wetlands are highly modified, and it receives significant discharges from agricultural drains and sewage effluent. The active floodplain is no longer functioning except in very high water conditions.”

Given the level of alteration, restoring the Lost River sub-basin would be a monumental task requiring conversion of thousands or tens of thousands of acres of farmland back to wetlands. This would require large amounts of money and political will which is unlikely to materialize. Therefore, the Consortium is not considering the Lost River watershed in this assessment to maintain focus on the main Klamath River and the greater possibility for restoration success. The problems of the Lost River can be addressed through treating the effluent prior to discharge into the Klamath River.

### 4.5 DAMS AND DIVERSIONS

Water levels in Upper Klamath Lake are regulated by Link Dam near Klamath Falls, Oregon. Downstream of this location, there are an additional five dams on the Klamath River that form the Klamath Hydroelectric Project: Keno (River Mile 233), J.C. Boyle (RM 225), Copco 1 (RM 198) and 2 (RM 196), and Iron Gate (RM 190) (Figure 2). Fish passage at Link Dam was recently upgraded, but fish ladders at Keno and J.C. Boyle dams do not meet current fish passage criteria and there are no fish passage facilities at Copco 1 & 2 and Iron Gate dams (U.S. DOI and CDFG 2012). Below Iron Gate Dam, the Klamath River is free flowing to the Pacific Ocean. Keno, J.C. Boyle, and Copco 1 & 2 and Iron Gate dams do not store water for irrigation.

The Klamath Hydroelectric Settlement Agreement (KHSA) was drafted in 2010 and signed by PacifiCorp (operator of the dams), the U.S. Secretary of the Interior, Oregon and California

governors, and multiple stakeholders. The KHSA called for the removal of Iron Gate, Copco 1 & 2, and J.C. Boyle dams, but it was never approved by the U.S. Congress. A revised KHSA was announced in 2016 which did not require congressional action and still calls for the removal of the lower 4 dams.

The U.S. Bureau of Reclamation Klamath Irrigation Project diverts water from Upper Klamath Lake, the Klamath River, and waterbodies in the Lost River basin for agricultural irrigation of >200,000 acres in California and Oregon. The Klamath Project has 19 major canals and laterals that form a 1200-mile-long irrigation network (USBR 2016a). There are additional diversions outside of the Klamath Irrigation Project, including both upstream and downstream of Upper Klamath Lake. The total amount of irrigation water consumed by crops in the Klamath Basin upstream of Iron Gate Dam including Lost River and Butte Valley is approximately 766 million cubic meters (621,000 acre-feet) per year (Asarian and Walker 2016).

#### 4.6 LAND USE AND OWNERSHIP

Land use in the Upper Klamath basin is predominantly public and private forestry. Agriculture and rangeland comprise only a small portion of the basin, but these activities are located in relatively sensitive areas of the basin, including valley bottoms and along the shores of UKL. In the Upper Klamath River watershed, large areas of wetlands have been drained, streams diked, and water diverted to support agricultural land use, which has resulted in increased concentrations of nutrients and sediment delivered to watercourses (Stillwater Sciences et. al 2012). A small portion of the upper basin is protected in Crater Lake National Park, Lava Beds National Monument, and in National Wildlife Refuges (ODEQ 2002).

Table 2. Land cover types in Upper Klamath Basin NPS AMPP area.

Land Cover Type	Square Kilometers	Square Miles	Percent of Total
Forest	12977.1	5010.5	54.5
Shrub/Scrub	4755.4	1836.1	20.0
Grassland/Herbaceous	2176.4	840.3	9.1
Cultivated Crops, Pasture/Hay	1567.6	605.3	6.6
Wetlands	1134.9	438.2	4.8
Open water	731.9	282.6	3.1
Developed	298.1	115.1	1.3
Barren (Rock/Sand/Clay)	189.9	73.3	0.8
Total	23831.3	9201.3	100.0




Figure 6. Land cover in the Upper Klamath basin. Data are from the Multi-Resolution Land Characteristics Consortium for the year 2011.




Figure 7. Map of land ownership in the Upper Klamath Basin. The dotted red line is the outline of the NPS project area. Map adapted from Stillwater Sciences et al. (2012).

#### 4.6.1 NATIONAL WILDLIFE REFUGES

The USFWS manages three National Wildlife Refuges (NWRs) in the Upper Klamath Basin. The Klamath Marsh NWR contains 41,230 acres of wet meadow and open-water wetland habitat located along the Williamson River. The Upper Klamath NWR protects 15,000 acres of emergent marsh and open water on the west shores of UKL. Lastly, the Lower Klamath NWR contains 50,092 acres of shallow marsh, open water, grassy upland, and cropland located southeast of the Keno Reservoir and the Klamath River. Of these three NWRs, the Lower Klamath NWR hydrology is the most impacted by allocation of water and in some years is unable to keep marshes wet.

#### 4.6.2 FORESTRY

Residual impacts from past harvesting practices in the Upper Klamath River include abandoned or non-maintained roads that have not been fully decommissioned, single-aged tree stands reforesting clearcut areas, and vegetation removal. These activities modify groundwater and surface hydrology and can contribute high loads of sediment and associated nutrients to streams. The Freemont-Winema National Forest (NF), Klamath NF, and Modoc NF all have active timber management areas within the Upper Klamath Basin. Private timber companies also own and manage land for timber harvest.

#### 4.6.3 AGRICULTURE

Agricultural activities in the Upper Klamath River area include cattle grazing and crop production. The majority of agricultural areas are located along the relatively flat valley bottoms near current or historic rivers, lakes, and wetlands. Due to the lack of summer precipitation, irrigation greatly increases agricultural productivity and therefore most agricultural lands are irrigated. Crops grown within the Upper Klamath Basin include cereals (barley, oats, and wheat), forage (alfalfa, hay, and irrigated pasture), and other crops (potatoes, sugar beets, onions, peppermint, horseradish, and pea seed) (Smith and Rykbost 2000). In the tributaries upstream of Upper Klamath Lake which is the primary focus of this NPS AMPP, irrigated pasture for cattle grazing predominates (NRCS 2009, 2010).

#### 4.6.4 URBAN

Klamath Falls is the largest metropolitan area in the Upper Klamath Basin (Figure 2). Located at the outlet of Upper Klamath Lake, Klamath Falls covers 20.6 mi<sup>2</sup> and has a population of 21,207. Other towns in the area include Chiloquin and the unincorporated communities of Ady, Beatty, Bly, Klamath Agency, and Worden.



## 5 WATER QUALITY SUMMARY

### 5.1 SYNTHESIS OF UPPER KLAMATH BASIN WATER QUALITY IMPAIRMENTS AND LINKS TO NPS POLLUTION

Upper Klamath Lake has historically been a eutrophic lake, but land-use practices and hydrologic modifications in the past several decades have caused the lake to become hyper-eutrophic (ODEQ 2002). There is widespread agreement that excessive levels of phosphorus are the ultimate cause of the pH and DO impairment in Upper Klamath Lake (ODEQ 2002). The Upper Klamath Lake TMDL (ODEQ 2002) presents details of research that identified sources of phosphorus and the pathway of phosphorus-induced impairment of pH and DO in UKL.

Conversion of wetlands to agricultural land has increased the loading of phosphorus into Upper Klamath Lake. Phosphorus is contributed to Upper Klamath Lake when wetlands are drained, and their peat soils are exposed to oxygen which allows decomposition and leads to release of phosphorus which is then delivered to the lake through runoff or pumping. The reclaimed wetlands are then used for cattle grazing and crop production. Runoff from these land uses is high in nutrients, including phosphorus.




Figure 8. Seasonal nutrient mechanisms in Upper Klamath Lake at a watershed scale. Figure copied from Stillwater Sciences et al. (2013).

High phosphorus loading and naturally warm water temperatures trigger cyanobacteria blooms in Upper Klamath Lake that themselves result in low DO and high pH in not only Upper Klamath Lake but also in reservoirs downstream. Cyanobacteria blooms in Upper Klamath Lake are severe and can occur simultaneously throughout the entire lake. The senescence of the blooms results in anoxic environments as the cyanobacteria decompose. Dead cyanobacteria settle into the sediments, providing an internal source of phosphorus which can fuel blooms later that year and the following year (Figure 8). Cyanobacteria-rich waters flow out of Upper Klamath Lake through Link River and into Lake Ewauna and Keno Reservoir (Figure 9). Turbulent conditions in Link River degrade the cyanobacteria, causing much of it to die (Sullivan et al. 2011). The dead cyanobacteria settle to the bottom of Keno Reservoir and consume oxygen as decomposition occurs (Sullivan et al. 2013, 2014).




Figure 9. Effect of Upper Klamath Lake algal blooms on Keno Reservoir and the Klamath River downstream. Figure copied from Stillwater Sciences et al. (2013).

## 5.2 PHOSPHORUS

Primary external sources of phosphorus in the Upper Klamath Lake watershed include decomposing peat in drained wetlands, springs transporting phosphorus dissolved from geologic sources of volcanic origin, and agricultural runoff. External loads of phosphorus contributed to the lakes account for nearly 40% of phosphorus concentrations on an annual basis (ODEQ 2002). The Williamson, Sprague and Wood rivers contributed 65% combined external phosphorus load from nearly 85% of the watershed area (Figure 10). ODEQ recommends a 40% reduction in total external phosphorus loading to Upper Klamath Lake as a target condition for the TMDL (ODEQ 2002).

Drained wetlands contribute large amounts of nutrients to Upper Klamath Lake. When intact and functioning, wetlands are a nutrient sink. The water that drains from reclaimed wetlands contains high concentrations of nutrients, including phosphorus and nitrogen (Snyder and Morace 1997, Kann and Walker 1999, ODEQ 2002). As wetlands dry after draining, aerobic peat decomposition releases additional nutrients into the waters pumped from the reclaimed areas (Snyder and Morace 1997, ODEQ 2002). Wetlands are most often drained for agricultural use, which includes cattle grazing and cultivation of crops (ODEQ 2002). Not only have the majority of wetlands around Upper Klamath Lake and Agency Lake been drained, so have wetlands in the Williamson and Sprague watersheds. Following the recognition of negative effects of wetland draining on habitat and water quality, significant progress has been made in the past two decades

to begin the process of restoring lakeside wetlands in the Williamson River Delta (Wong and Hendrickson 2011) and the mouth of the Wood River (Carpenter et al. 2009). The percent of the lake's total external phosphorus load that is attributable to anthropogenic land-use activities declined from 38% (1992–1998) to 31% (2008–2010), in part due to wetland restoration which decreased the amount of drainage water pumped from agricultural lands into the lake (Walker et al. 2012).

Other human activities and land uses besides wetland conversion have also increased external loads of phosphorus to Upper Klamath Lake. A primary mechanism appears to be increased rates of erosion and transport of sediment-bound phosphorus (ODEQ 2002, Walker et al. 2015). For example, sediment cores within Upper Klamath Lake indicate an increase in various indicators (e.g. Ti, Al, tephra, and charcoal) of erosional inputs in the 20th century (Bradbury et al. 2004, Eilers et al. 2004, Simon and Ingle 2011). The Sprague River is an important source of particulate-bound phosphorus (ODEQ 2002, Walker et al. 2015). Woody vegetation has decreased in the past century in the valley bottoms along the Sprague River and Sycan River (O'Connor et al. 2015). Levees confine stream channels in parts of the Sprague River watershed, disconnecting floodplains and promoting incision which results in sediment being transported downstream rather than being deposited on floodplains (O'Connor et al. 2015). Monitoring of pastures in the Upper Klamath Lake basin found that large quantities of phosphorus can be exported into watercourses during first-flush irrigation events and storm events (Ciotti et al. 2010).

Recycled sediment and algal phosphorus (i.e., internal loading) in Upper Klamath Lake and Agency Lake contributes nearly 60% of total in-lake phosphorus concentrations on an annual basis (ODEQ 2002). Mechanisms for internal recycling vary spatially and temporally but likely include a combination of algal translocation, diffusion, pH or anaerobic-mediated release, microbial and macroinvertebrate metabolic cycling, bioturbation, and resuspension (Barbiero and Kann 1994; Laenen and LeTourneau 1996; Kuwabara et al. 2007, 2009, 2012; Simon et al. 2009; Simon and Ingle 2011). If external loads can be reduced, internally recycled phosphorus would likely reach a new lower equilibrium level within a few decades (ODEQ 2002, Wherry et al. 2015).




Figure 10. Relative contributions of tributaries and other sources to external total phosphorus (TP) load, inflow water volume, and drainage area to Upper Klamath Lake for hydrologic years 1992-2010. Figure adapted from Stillwater Sciences et al. (2012) based on data from Walker et al. (2012).

### 5.3 NITROGEN

*Aphanizomenon flos-aquae* is capable of fixing nitrogen. Mass-balance nutrient budgets for Upper Klamath Lake find net-negative retention of nitrogen during the summer bloom period, consistent with large quantities of N being fixed (Walker et al. 2012). N:P ratios and overall nitrogen concentrations coming into UKL are very low, favoring nitrogen-fixers such as *Aphanizomenon flos-aquae* which can overcome nitrogen limitation. Therefore, while nitrogen concentrations are monitored within the NPS AMPP area, the Consortium’s efforts focus on phosphorus because it is the most feasible means of controlling *Aphanizomenon flos-aquae* within Upper Klamath Lake. Controlling *Aphanizomenon flos-aquae* in UKL is critically important since its biomass sets the stage for poor water quality and blooms of toxic *Microcystis* both in UKL and downstream in the Klamath River.

### 5.4 DISSOLVED OXYGEN AND PH

Due to prolific blooms and crashes of the cyanobacterium *Aphanizomenon flos-aquae*, low dissolved oxygen (DO) and high pH conditions can occur in Upper Klamath Lake during late spring, summer, and early fall. Low DO, high pH, and high ammonia concentrations have been identified as primary factors in the declines of the endangered shortnose sucker (*Chasmistes brevirostris*) and Lost River sucker (*Deltistes luxatus*) in Upper Klamath Lake (ODEQ 2002). DO concentrations in the lake can drop below the ODEQ warmwater aquatic life criterion of 5.5 mg/L for many weeks in a row (Kann 2010, Morace 2007). Annual timing is variable depending on the bloom cycle, but pH peaks in July during blooms and DO is often lowest in August when blooms are declining and water temperatures are warm (Jassby and Kann 2010).

DO conditions in Lake Ewauna and Keno Reservoir are even more acute than in Upper Klamath Lake, approaching zero for weeks or months at a time (Sullivan et al. 2013, 2014).

The Klamath Hydroelectric Project (KHP) has a direct effect on DO and pH levels in the Klamath River immediately below Iron Gate Dam (FERC 2007). During the summer season the reservoir often releases water with high pH and low DO (Asarian and Kann 2013), which could harm salmonid fish in the vicinity of the dam. Phytoplankton blooms from KHP reservoirs tend to decrease daily minimum dissolved oxygen concentrations in the Klamath River, presumably by reducing light availability and rates of production from periphyton (Genzoli 2013, Genzoli et al. 2015, Genzoli and Hall 2016).

Downstream of our NPS project area in the free-flowing reaches of the Klamath River, photosynthesis and respiration by periphyton (algae attached to the riverbed) and aquatic plants in the Klamath River can degrade dissolved oxygen and pH conditions, resulting in water quality that is chronically stressful to fish (NCRWQCB 2010, Asarian and Kann 2013).

## 5.5 TEMPERATURE

High water temperatures in Upper Klamath Basin streams are due in part to solar radiation from poor riparian conditions and contributions of irrigation return flows that are warmer than ambient river flows (ODEQ 2002). Other hydrologic modifications in the UKL watershed include water diversions, reduction in stream flows, streambed channelization, streambed armoring, dikes, and dams.

Upper Klamath Lake's shallow depth prevents thermal stratification and results in naturally high summer water temperatures throughout much of the lake; however, springs do provide cold-water refugia for fish and other organisms in the lake.

Primarily due to the thermal mass of Iron Gate and Copco reservoirs, the Klamath Hydroelectric Project significantly alters water temperatures in the Klamath River (PacifiCorp 2004, 2005; FERC 2007, US DOI and CDFG 2012) in ways that are detrimental to the various runs of anadromous fish in the Klamath River.

## 5.6 CYANOBACTERIAL TOXINS

Iron Gate and Copco reservoirs provide ideal habitat for the toxigenic cyanobacteria (blue-green algae) *Microcystis aeruginosa* by transforming turbulent free-flowing river reaches into stagnant thermally stratified impoundments that favor cyanobacterial proliferation. In the presence of abundant nutrients, the transformation from river to reservoir environment leads to massive blooms (Kann 2006; Kann and Corum 2006, 2007). Microcystin toxins produced by the toxic cyanobacteria *Microcystis* represent a substantial threat to human and animal health (OEHHA 2005; Kann 2006; Kann and Corum 2006, 2007; OEHHA 2012). The Klamath River is listed as impaired by microcystin toxins from Stateline to its confluence with the Trinity River (SWRCB 2015). Microcystin concentrations generally decline with distance downstream of Iron Gate Dam (US DOI and CDFG 2012) but frequently exceed public health guidelines between Iron Gate and Orleans, and occasionally exceed public health and water quality criteria as far downstream as the Klamath Estuary (HVTEPA 2013). Genetic fingerprinting research showed that Iron Gate Reservoir is the source of downriver *Microcystis* assemblages and that Iron Gate Reservoir was determined to be the principal source of *Microcystis* found throughout the lower 300 km of river

separating the reservoir from the Pacific Ocean (Otten et al. 2015). Tissue sampling has documented bioaccumulation of microcystin in freshwater mussels from the Klamath River between Iron Gate Dam and the Yurok Reservation, yellow perch from Iron Gate and Copco reservoirs, juvenile salmonids in Iron Gate Hatchery, and steelhead and Chinook salmon in the Klamath River (Fetcho 2006, Kann 2008, OEHHA 2008, and Kann et al. 2010).

*Microcystis* and microcystin also occur upstream in Upper Klamath Lake (Kann et al. 2015), particularly in the northern part of the lake (Agency Lake), but typically at levels far lower than is observed in Iron Gate and Copco reservoirs (Kann 2006). For reasons that have not yet been determined, there were large blooms of *Microcystis* in Upper Klamath Lake in 2014 and 2015.

## 5.7 SEDIMENT

Several tributaries to the Klamath River are listed as sediment impaired on the 303(d) list (Appendix A). Land uses contributing to sediment to waterbodies include timber harvest and associated roads, cattle grazing, other agriculture, and degraded riparian conditions. Sediment cores in Upper Klamath Lake show strong evidence of increased erosion inputs in the lake in the past century relative to previous periods (see section 5.2 above). Increased sedimentation can directly impact aquatic resources within tributaries as well as contribute to phosphorus loading and algal blooms within Upper Klamath Lake.

# 6 RESULTS

As discussed above in Section 3, since the state of California and Oregon have already done a thorough analysis of impairment on individual waterbodies including the development of TMDLs (ODEQ 2002; 2010b, NCRWQCB 2010), we rely heavily on their existing assessments.

## 6.1 BENEFICIAL USES

Beneficial uses of the Klamath River and its tributaries and lakes in Oregon that have been identified by the Oregon Department of Environmental Quality (ODEQ) include direct human contact, agricultural, and aquatic life uses (Table 3). In California, designated beneficial uses are listed in Table 4 for the Middle Klamath River from Hornbrook Creek upstream to Copco Reservoir, not including tributaries.

Water bodies in California and Oregon are assessed and placed in a reporting category that represents levels of water quality attainment and support of beneficial uses. These categories (Table 5) are developed according to U.S. EPA guidance for Integrated Reports (U.S. EPA 2005). The categories range from Category 1, where all beneficial uses are fully supported, to Category 5, where at least one beneficial use is not supported and a TMDL is required.

A beneficial use is considered not supported if a water body does not meet designated water quality standards. The standards are set by each state and approved by USEPA. Some standards are numeric (i.e., pH, dissolved oxygen, chlorophyll-a) while others are narrative, such as “no measurable surface water temperature increase resulting from anthropogenic activities is allowed” (ODEQ 2002). For example, ODEQ states the beneficial use for salmonid fish rearing is considered fully supported if water temperatures do not exceed 17.8°C (ODEQ 2002).

The majority of the beneficial uses in Oregon are impaired by parameters from non-point sources, including chlorophyll-a, pH, and temperature. In California, the Klamath River does not

fully support the following beneficial uses that include cold freshwater habitat; rare, threatened and endangered species; migration of aquatic organisms; spawning, reproduction, and early development of fish; commercial and sport fishing; Native American cultural use; subsistence fishing; and contact and non-contact water recreation (NCRWQCB 2010).

Table 3. ODEQ Designated Beneficial Uses for the Upper Klamath River and other Basin waters in Oregon. The other waters include Williamson, Sprague, and Sycan watersheds and tributaries and lakes below Keno Dam. Table modified from ODEQ Designated Beneficial Uses Klamath Basin (340-41-0180) Table 180A (ODEQ 2005).

<b>Designated Beneficial Use</b>	<b>Klamath River (Upper Klamath Lake to Keno Dam)</b>	<b>All Other Basin Waters</b>
Domestic Water Supply	x	x
Industrial Water Supply	x	x
Hydropower	x	
Water Contact: Recreation	x	x
Irrigation	x	x
Livestock Watering	x	x
Wildlife and Hunting	x	x
Fishing	x	x
Commercial Navigation and Transportation	x	
Boating	x	x
Aesthetic Quality	x	x
Resident Fish and Aquatic Life	x	x
Salmonid Fish Rearing	x	
Salmonid Fish Spawning	x	
Anadromous Fish Passage	x	
Coldwater Fisheries	x	x

Table 4. NCRWQCB Designated Beneficial Uses for the Middle Klamath River from Hornbrook upstream to Copco Lake. Data modified from Water Quality Control Plan for the North Coast Region Table 2.1 (NCRWQCB 2011). E: Existing designated beneficial use; P: Potential beneficial use.

<b>Designated Beneficial use</b>	<b>Hornbrook to Iron Gate Dam</b>	<b>Iron Gate Reservoir</b>	<b>Copco Lake</b>
Municipal and Domestic Supply	E	P	E
Agricultural Supply	E	P	E
Industrial Service Supply	E	P	E
Industrial Process Supply	E	P	P
Groundwater Recharge	E		
Freshwater Replenishment	E	E	E
Navigation	E	E	E
Hydropower Generation	P	E	E
Water Contact Recreation	E	E	E
Non-Contact Water Recreation	E	E	E
Commercial and Sport Fishing	E	E	E
Aquaculture	P		
Warm Freshwater Habitat	E	E	E
Cold Freshwater Habitat	E	E	E
Wildlife Habitat	E	E	E
Rare, Threatened, or Endangered Species	E	E	E
Migration of Aquatic Organisms	E	E	E
Spawning, Reproduction, and/or Early Development	E	E	E
Shellfish		E	

Table 5. Assessment categories of water body support of designated beneficial uses. Categories are based upon USEPA guidance (2005) and utilized by California (SWRCB 2015b) and Oregon (ODEQ 2012). Category 3b is used by Oregon only.

<b>Category</b>	<b>Definition</b>
1	A water segment that supports all core beneficial uses.
2	A water segment that has some core beneficial uses supported.
3	Evidence is insufficient to make a supportive beneficial use determination.
3b	Potential concern, data are insufficient to determine beneficial use support.
4a	A water segment that has at least one beneficial use not supported and has a U.S. EPA-approved TMDL
4b	A water segment that has at least one beneficial use not supported and a TMDL is not required as an existing regulatory program addresses the issue.
5	A water segment that has at least one beneficial use not supported and a TMDL is required but not yet completed.



## 6.2 WATER QUALITY LIMITED WATERS

ODEQ has listed over 45 stream segments and six reservoirs and lakes as 303(d) water quality limited and either requiring a TMDL or a TMDL has been approved. The list was submitted by ODEQ to EPA in May 2011 and approved by EPA in March 2012. See Appendix A for the 2012 ODEQ impaired waters list for streams, lakes, and reservoirs in the Klamath Basin that are within Oregon state boundaries.

Some sections of rivers, streams, and lakes are not 303(d) listed as impaired due to either insufficient data or the pollutant is potentially of concern for the stream section. Some waterbodies are considered water quality limited but a TMDL is not required due to the pollutant not being regulated by TMDLs, including flow and habitat modifications. Phosphorus is a pollutant of major concern in the Upper Klamath Basin. The ODEQ TMDL (ODEQ 2002) for Upper Klamath Lake describes a clear relationship between phosphorus concentrations and the listed impairments of pH, dissolved oxygen, and chlorophyll-a, a finding corroborated by many other analyses (Kann 1993, 1998; Wherry et al. 2015). If phosphorus loading to Upper Klamath Lake and Agency Lake can be reduced, it is highly likely that the magnitude of algal blooms in Upper Klamath Lake would also be reduced (ODEQ 2002, Wherry et al. 2015), which would improve water quality of the Klamath River and its reservoirs downstream (NCRWQCB 2010, Sullivan et al. 2013).

The North Coast Regional Water Quality Control Board in California has designated the Klamath River as “impaired” for water temperature, dissolved oxygen (DO), organic matter (measured as Carbonaceous Biochemical Oxygen Demand, CBOD), total phosphorus (TP), total nitrogen (TN), and microcystin (NCRWQCB 2010). The NCRWQCB TMDL for the Klamath River (2010) states in reference to waters entering from Oregon that “current source loads have overwhelmed the historical renewal capabilities of the Klamath River, leading to its impaired status.” Current annual loads of TP, TN, and CBOD in the Klamath River at Stateline (the Oregon-California border) are more than twice as high as estimated natural conditions (NCRWQCB 2010).

### 6.3 ATTRIBUTION OF IMPAIRMENTS TO NPS POLLUTION FOR EACH SUB-BASIN

Table 6. Sub-basin summary of impairments and causes of NPS pollution. For the sake of simplicity and usefulness, the table does not list every impairment for every specific waterbody but rather focuses on the most important impairments and causes at the sub-basin level. Waterbodies are listed separately in the Hydroelectric Reach because the waterbodies have different causes of impairment.

<b>Sub-basin</b> <i>Waterbody</i>	<b>Impaired Parameter</b>	<b>Severity</b>	<b>NPS Pollution Categories [i.e., Causes] and Importance</b>		
<b>Williamson River</b>	Phosphorus	Moderate	<i>Primary:</i> Agriculture (AGR), Hydromodification and Habitat Alteration (HHA)		
	Temperature	Moderate			
<b>Sprague River</b>	Phosphorus	High	<i>Primary:</i> Agriculture (AGR), Hydromodification and Habitat Alteration (HHA)		
	Dissolved Oxygen	Moderate			
	pH	Moderate			
	Temperature	Moderate			
<b>Wood River and other tributaries to UKL</b>	Phosphorus	High	<i>Primary:</i> Agriculture (AGR), Hydromodification and Habitat Alteration (HHA)		
<b>Upper Klamath Lake including Agency Lake</b>	Phosphorus	High	<i>Primary:</i> Agriculture (AGR), Hydromodification and Habitat Alteration (HHA)		
	Algae	High			
	Chlorophyll-a	High			
	Dissolved Oxygen	High			
	pH	High			
	Ammonia	High			
<b>Link River, Lake Ewauna, Keno Reservoir, and Lower Klamath Lake</b>	Phosphorus	High	<i>Primary:</i> Agriculture (AGR), Hydromodification and Habitat Alteration (HHA)  <i>Secondary/Minor:</i> Urban (URB)		
	Algae	High			
	Chlorophyll-a	High			
	Dissolved Oxygen	High			
	pH	High			
	Ammonia	High			
	Temperature	Moderate			
<b>Hydroelectric Reach (Keno Dam to Iron Gate Dam)</b> <i>Mainstem River and Reservoirs</i>	Cyanobacterial toxins	High	<i>Primary:</i> Hydromodification and Habitat Alteration (HHA), Agriculture (AGR)		
	Algae	High			
	Chlorophyll-a	High			
	Dissolved Oxygen	Moderate			
	pH	Moderate			
	Temperature	Moderate			
	<i>Spencer Creek</i>	Sedimentation		Moderate	<i>Primary:</i> Forestry (FOR), Agriculture (AGR), Hydromodification and Habitat Alteration (HHA)
		Temperature		Moderate	

## 7 DISCUSSION

Linkages between nonpoint source pollution and water quality problems in the Upper Klamath Basin have been intensively studied for decades and are relatively well understood. Human activities have pushed the naturally eutrophic Upper Klamath Lake over the edge into a hyper-eutrophic state. Excessive loading of phosphorus into the lake has fueled prolific blooms of the nitrogen-fixing cyanobacterium *Aphanizomenon flos-aquae* which then severely degrade water quality conditions both within the lake as well as downstream when the algae decay, consume dissolved oxygen, and release ammonia.

Simply put, the key to reversing the problem is to find ways to keep phosphorus on the land and prevent it from entering Upper Klamath Lake and its tributaries. This should be the highest priority. A secondary strategy is to attempt to remove phosphorus that has already reached the tributaries, such as by routing water through treatment wetlands or restoring floodplain connectivity so that sediment-bound phosphorus deposits on floodplains rather than being transported downstream into the lake. A tertiary strategy, which is less desirable because it will only help the river downstream, not the lake itself, would be to intercept (with mechanical removal or treatment wetlands) algae at the outlet of the lake before it is discharged to the Klamath River downstream. A fourth strategy would be to mechanically oxygenate Keno Reservoir, which is the location of some of the worst water quality in the entire Klamath Basin and will be a key bottleneck for recolonization of salmon into the Upper Klamath Basin once the dams are removed.

Human activities contributing phosphorus loads to Upper Klamath Lake include conversion wetlands to agricultural land, pumping of water from former wetlands directly into the lake or its tributaries, suppression of riparian vegetation by cattle grazing or mechanical disturbance, discharge of cattle manure into streams, return of irrigation tailwater to streams, construction of levees which confine stream channels and promote sediment transport rather than sediment retention, and roads and timber harvest which cause erosion in upland areas.

Cyanobacteria, including the toxigenic *Microcystis aeruginosa*, also bloom in the hydroelectric reservoirs downstream of Keno Dam. The most effective means for preventing these blooms is to remove the dams, as has been proposed, and convert the warm quiescent waters of the reservoirs back to turbulent free-flowing river reaches.

## 8 SELECTION OF BEST MANAGEMENT PRACTICES

### 8.1 CORE PARTICIPANTS

NPS pollution prevention is the responsibility of all those who live and work in the Upper Klamath Basin. As such, cooperation between all entities is vital to the health of the watershed.

The Consortium will combine sound science and local knowledge to decide which BMPs (see below) to support. It is not the Consortium's intention to initiate BMPs or restoration projects within the Upper Klamath Basin, but rather to support organizations and programs that already implement effective BMPs and projects that provide a positive impact to water quality. As part

of that support, the Consortium will continue to develop working relationships with multiple federal and tribal agencies operating within the Upper Klamath Basin on water quality issues.

Table 7 identifies the institutions that are involved in the identification of NPS pollution and impaired waterways, selection and approval of BMP's for NPS pollution, implementation of NPS pollution BMPs and all other key planning and management documents. Additional discussion for each of these agencies is included in the Consortium's Nonpoint Source Pollution Management Program Plan (Section 11) and/or Section 9.

Table 7. Core participants for BMPs

<b>Entity Category</b>	<b>Participant</b>	<b>Role</b>
<b>Tribal</b>	Klamath Tribal Water Quality Consortium	Collaborates with entities who are working to restore water quality in the Upper Klamath Basin. Provides technical and financial support for projects to improve water quality.
	Quartz Valley Indian Reservation, Yurok Tribe, Karuk Tribe, Hoopa Tribe, and Resighini Rancheria	Member Tribes of the Klamath Tribal Water Quality Consortium.
	Klamath Tribes of Oregon	Conducts water quality monitoring, plans/implements projects to improve habitat and water quality. Ancestral territory spans large portion of the Upper Klamath Basin.
<b>Federal</b>	US Environmental Protection Agency	Provides regulatory and technical assistance as well as grant funding for tribal environmental projects. Reviews and approves the Consortium's NPS Assessment Report and Management Plan.
	U.S. Bureau of Reclamation	Operates the Klamath Irrigation Project and provides funding for research, monitoring, and habitat restoration.
	U.S. Fish and Wildlife Service	Provides funds for habitat restoration on private and tribal lands. Manages several National Wildlife Refuges.
	U.S. Bureau of Land Management (U.S. BLM)	Manages federal land including forests, rangelands, and the Wood River Wetland.
	U.S. Forest Service (USFS)	Manages federal land including forests and rangelands
	Natural Resources Conservation Service (NRCS)	Provides funding, technical oversight, and education for projects on private agricultural lands.
	National Oceanic and Atmospheric Administration (NOAA)	Provides funding under the Congressionally mandated Pacific Coastal Salmon Recovery Fund (PCSRF).
<b>State</b>	California State Water Resources Control Board (SWRCB) and North Coast Regional Water Quality Control Board (NCRWQCB)	Regulates implementation of California Nonpoint Source Plan, the North Coast Basin Plan, Waste Discharge Requirements, and Total Maximum Daily Loads. Promotes stewardship and collaboration.
	California Department of Fish and Wildlife	Provides funding for instream and upslope restoration projects. Provide permits for projects that require alteration of streambanks.
	California Coastal Conservancy	Provides funding for habitat restoration projects.

Entity Category	Participant	Role
	Oregon Department of Environmental Quality (ODEQ)	Regulates water quality in the State of Oregon.
	Oregon Department of Agriculture (ODA)	Designated Management Agency which develops implementations plan for addressing nonpoint source pollution from agricultural lands. Provides education and technical assistance to farmers and ranchers.
	Oregon Watershed Enhancement Board (OWEB)	Provides funding for habitat restoration projects.
<b>Other Government</b>		
	Klamath Soil and Water Conservation District (KSWCD)	Provides education and technical assistance to farmers and ranchers.
	Klamath Basin Research & Extension Center (KBREC)	Provides education and technical assistance to farmers and ranchers.
<b>Private Companies</b>		
	PacifiCorp	Operates the Klamath Hydroelectric Project and provides funding for water quality projects under the Klamath Hydroelectric Settlement Agreement (KHSa).
<b>Non-Profit Organizations</b>		
	Klamath Watershed Partnership (KWP)	Provides outreach, education, and technical assistance to residents and agricultural operators. Implements projects to improve habitat and agricultural operations.
	The Nature Conservancy (TNC)	Acquires, manages, and restores land with high conservation value, including wetlands.
	Trout Unlimited/Klamath Basin Rangeland Trust (TU/KBRT)	Implements projects to improve water quality, water quantity, and fish habitat.
<b>Other</b>		
	Klamath Basin Monitoring Program (KBMP)	Coordinates basin-wide water quality monitoring and facilitates collaboration and information sharing.

## 8.2 PUBLIC PARTICIPATION AND GOVERNMENTAL COORDINATION

NPS pollution is a community-wide issue and successful implementation of the NPS AMPP will rely upon relationships between the Consortium, our partner entities, and the public, including tribal and non-tribal community members. Therefore, the Consortium sought public input on this NPS AMPP by engaging public agencies that have a role in managing or protecting natural resources. The Consortium did an oral presentation on the NPS AMPP at the spring 2016 Klamath Basin Monitoring Program (KBMP) meeting which was attended by approximately 50 people, primarily representatives of entities involved in Klamath Basin water quality issues. The Consortium and its consultants conducted phone meetings with many partner organizations to get input on the NPS AMPP, and followed up with email correspondence.

In addition, the Consortium made a draft version of this document available for a 30-day public comment period starting August 19, 2016. Public notice was made by announcing the release of the document in each Consortium Tribe's newsletter, official website, and social media sites. The public notice was also listed on the KBMP website. The Consortium will review all comments

that were received and consider them thoroughly before making appropriate changes to the document.

## 9 EXISTING NPS CONTROL PROGRAMS

There are many groups Upper Klamath Basin working to reduce nonpoint source pollution within the Upper Klamath Basin. This section describes some of the active organizations and the types of projects implemented.

### 9.1 FEDERAL AGENCIES

#### 9.1.1 U.S. BUREAU OF RECLAMATION

As noted in Section 4.5 above, the U.S. Bureau of Reclamation (USBR) operates the Klamath Irrigation Project (KIP). USBR has several programs relevant to NPS control.

Under the Klamath Basin Water Supply Enhancement Act of 2000<sup>3</sup>, USBR is authorized to conduct feasibility studies on ways to improve water supply, water quality, and fish and wildlife habitat. That source of funds is limited to reconnaissance, appraisal, and feasibility, and may not be used for implementation. USBR recently completed an initial assessment of potential projects to improve water quality and supply within the KIP (USBR 2016b). The assessment identified 39 site-specific projects which included treatment wetlands, water reuse, water recirculation, and water storage (USBR 2016b). USBR is now planning to further evaluate a subset of the projects included in the initial assessment (Rick Carlson, personal communication, June 17, 2016).

The Recovery Implementation Team (RIT) for Lost River and shortnose suckers was formed following the revision of the recovery plan for those species (USFWS 2012) and is funded by USBR. The RIT issues solicitations for projects and selects projects for funding. The projects typically focus on sucker biology and ecology but some also include linkages to water quality. Examples of previously funded projects included an ongoing adult sucker population monitoring program, lab studies on sucker tolerance for microcystin toxins, juvenile sucker cohort studies to track recruitment, and quantification of avian predation.

USBR also funds ongoing water quality monitoring, research, and analyses within UKL (Kann 1998, Lindenberg et al. 2009, Eldridge et al. 2014, Wood et al. 2013, Kann et al. 2015), and nutrient loading from lake tributaries (Kann and Walker 1999, Walker et al. 2012, Walker et al. 2015).

#### 9.1.2 U.S. FISH AND WILDLIFE SERVICE

The U.S. Fish and Wildlife Service (USFWS) Partners for Fish and Wildlife program provides funds for habitat restoration on private and tribal lands (USFWS 2013a). The Klamath Falls office of the USFWS coordinates the Partners for Fish and Wildlife projects within the Upper Klamath Basin. Potential projects include stream channel restoration, fish passage, riparian fencing and planting, wetland restoration, and spring reconnection. Although the goal of these projects is for fisheries and wildlife habitat restoration, many projects will have ancillary benefits to water quality. USFWS manages several national wildlife refuges within the Consortium's

<sup>3</sup> <https://www.govtrack.us/congress/bills/106/s2882>

NPS AMPP area. Along with its partners, the USFWS developed an innovative Walking Wetlands program on some refuge lands which involves a four-year rotation of crop production and inundation to create wetlands, resulting in nutrient retention, reduced weeds, wildlife habitat and improved crop production (USFWS 2013b).

### 9.1.3 NATURAL RESOURCES CONSERVATION SERVICE

The U.S. Department of Agriculture's (USDA) Natural Resources Conservation Service (NRCS) provides technical assistance in conservation planning concerning on-farm conservation implementation through federally funded programs supported by the Farm Bill. Through these programs, landowners can install fencing and watering systems to prevent cattle from trampling creeks and riparian areas, obtain subsidies for land put in conservation easement, implement irrigation improvements, build diffuse-treatment wetlands, and implement grazing rotational practices.

### 9.1.4 U.S. BUREAU OF LAND MANAGEMENT AND U.S. FOREST SERVICE

The U.S. Bureau of Land Management (BLM) manages forests and rangelands scattered around the Klamath Basin. Within the Consortium's NPS AMPP area, the BLM's two most relevant holdings are upland areas between Keno Dam and Iron Gate Dam (Figure 3), and the Wood River Wetland at the mouth of the Wood River. BLM is currently restoring the wetland habitats in Wood River Wetland (Carpenter et al. 2009). The U.S. Forest Service owns much of the upland forests in the Consortium's NPS AMPP area (Figure 3).

USFS and BLM forest lands are managed according to the Northwest Forest Plan (NWFP), a comprehensive federal ecosystem management strategy (USFS and BLM 1994). A primary feature of the NWFP is the Aquatic Conservation Strategy which includes watershed analyses, watershed restoration, and designation of riparian reserves and key watersheds.

### 9.1.5 U.S. ENVIRONMENTAL PROTECTION AGENCY

In collaboration within the Oregon Department of Environmental Quality and California's North Coast Regional Water Quality Control Board, the U.S. Environmental Protection Agency (U.S. EPA) developed and approved Total Maximum Daily Loads (TMDLs) specifying required water quality improvements within the Klamath and Lost rivers (U.S. EPA 2008, ODEQ 2010b, NCRWQCB 2010). U.S. EPA is actively involved in monitoring harmful algal blooms within the Klamath River and reservoirs, including processing cyanotoxin samples in its laboratory (Watercourse Engineering 2015).

## 9.2 THE KLAMATH TRIBES OF OREGON

The Klamath Tribes have lived in the Upper Klamath Basin since time immemorial. Following colonization by the United States and the resulting conflict, the Klamath Tribes signed a treaty in 1864 with the United States ceding much of their territory in exchange for a reservation and hunting and fishing rights. In 1954, federal recognition of the Klamath Tribes was terminated by the United States Congress. Federal recognition was restored in 1986 but the reservation was not returned (Klamath Tribes 2015).

The Klamath Tribes' Natural Resource Department initiated an ongoing long-term water quality monitoring program for Upper Klamath Lake and its tributaries in 1990 (Kann 1998, Kann and

Walker 1999, Walker et al. 2012). They also play a lead role in restoration planning in the Upper Klamath Basin.

### 9.2.1 WATER RIGHTS AND THE UPPER KLAMATH BASIN COMPREHENSIVE AGREEMENT (UKBCA)

The Oregon Water Resources Department (OWRD) began an administrative process for determining pre-1909 water rights in 1975, culminating in a Findings of Fact and Order of Determination (FFOD) in 2013 that the Klamath Tribes' water rights to Upper Klamath Lake and instream flows in the lake's tributaries dated to time immemorial. The Klamath County Circuit Court is now making the final adjudication, but while that process is pending the Klamath Tribes can enforce their water rights by making "calls" to OWRD to shut off diversions by junior water users (including some dating back as far as the 1860s) when instream flows are not met (Timmons 2016).

In response to the prospect of regular curtailment of irrigation diversions to meet tribal water rights, stakeholders upstream of Upper Klamath Lake negotiated the Upper Klamath Basin Comprehensive Agreement (UKBCA) which was signed in 2014 (UKBCA 2014). Signatories to the UKBCA include the Klamath Tribes, federal agencies, the State of Oregon, and water users. In exchange for implementation of the UKBCA's Water Use Program (WUP), Riparian Program, and Tribal economic development program, the Klamath Tribes agreed to not enforce their full water rights. The WUP mandates that instream flows into Upper Klamath Lake be increased by 30,000 acre-feet per year and establishes Specified Instream Flows (SIFs) for specific lake tributaries. The SIFs vary monthly according to hydrologic conditions. The purpose of the Riparian Program is to manage and restore riparian corridors along streams that flow into Upper Klamath Lake in order to achieve Proper Functioning Conditions. The Riparian Program establishes Riparian Management Areas along streams, which vary in width from 30 feet to 130 feet depending on site-specific conditions. The signatories of the UKBCA will jointly establish an Upper Klamath Basin Riparian Management and Restoration Action Plan (the Riparian Action Plan). The Riparian Action Plan will include parcel-specific management and restoration actions. These specific actions will then be incorporated into a Riparian Management Agreement for each property within the Riparian Management Corridor. The Klamath Tribes are currently negotiating the details of the Riparian Action Plan with the other stakeholders, with a targeted completion date of summer 2016 (Megan Skinner, personal communication, June 24, 2016). By 2018, 80% of the length of the Riparian Management Corridor must be enrolled in the Riparian Program or the Klamath Tribes will enforce their full water rights. Landowners will be responsible for maintenance but not initial building of fences. The Klamath Tribes and their partners are currently seeking funding to implement the UKBCA.

### 9.2.2 WATERSHED RESTORATION PLAN

Outside of the UKBCA process, the Klamath Tribes are also developing a Watershed Restoration Plan for the tributaries of Upper Klamath Lake which will include lands outside of the Riparian Management Corridor. The target date for completing the plan is the end of 2018.



## **9.3 STATE AGENCIES**

### **9.3.1 OREGON DEPARTMENT OF ENVIRONMENTAL QUALITY AND OREGON DEPARTMENT OF AGRICULTURE**

Oregon Department of Environmental Quality (ODEQ) developed Total Maximum Daily Loads (TMDLs) for two areas within the Consortium’s NPS project area, first Upper Klamath Lake and its tributaries (ODEQ 2002) and then the Oregon portion of the Klamath and Lost River basins (ODEQ 2010b). The TMDLs specify the Oregon Department of Agriculture (ODA) as the “Designated Management Agency (DMA)” which must develop an implementation plan for addressing nonpoint source pollution from agricultural lands. Since adoption of the TMDLs, ODA has developed agricultural water quality management plans for Klamath Headwater and Lost River sub-basins in collaboration with local agricultural water quality advisory committees and the Klamath Soil and Water Conservation District. The plans were first adopted in 2004 and have been subsequently revised (ODA 2011, ODA and LRLAWQAC 2015). The plans rely heavily on voluntary measures.

### **9.3.2 CALIFORNIA STATE WATER RESOURCES CONTROL BOARD AND NORTH COAST REGIONAL WATER QUALITY CONTROL BOARD**

The North Coast Regional Water Quality Control Board (NCRWQCB) and its related agency the California State Water Resources Control Board (SWRCB) have developed TMDLs throughout the California portion of the Klamath Basin, including the Klamath River TMDLs within the Consortium’s NPS project area (NCRWQCB 2010).

The California NPS Plan (SWRCB 2015a) describes California’s approach to addressing NPS pollution and designates the Klamath River as a statewide NPS priority. The Plan says California will continue with its Watershed Stewardship Approach in the Klamath River, which emphasizes coordination, collaboration, and building partnerships among entities. Another important component listed in the Plan is the updating of the Waiver of Waste Discharge Requirements (WDR) for federal land management activities within the North Coast Region. This federal waiver was renewed in 2015 and attempts to reduce water quality impacts from the following activities: timber harvesting, roads, grazing, recreation, vegetation management, restoration, fire suppression, and fire recovery (NCRWQCB 2015).

## **9.4 PRIVATE COMPANIES**

### **9.4.1 PACIFICORP**

PacifiCorp is a private company that operates the Klamath Hydroelectric Project (see Section 4.5 above). As part of the Klamath Hydroelectric Settlement Agreement (KHSA), PacifiCorp agreed to fund a number of Interim Measures (IM) including \$500,000 per year on water quality monitoring and \$250,000 to evaluate approaches for improving water quality including pilot studies (PacifiCorp 2014). The Interim Measures Implementation Committee (IMIC) is an interagency group that advises PacifiCorp on its IM study plans. Two members of the Consortium (the Yurok Tribe and Karuk Tribe) are signatories to the KHSA and are active participants in the IMIC. Examples of projects evaluated in IM studies include reducing instream nutrient loads using treatment wetlands (Lyon et al. 2009; CH2M HILL 2012, 2014) as well as removal of algal biomass near Link Dam using stormwater technology (hydrodynamic separators, Watercourse Engineering, Inc. 2013, 2014a, 2014b) or other methods (PacifiCorp

2015, Carlson and Hughes 2016). Other IM studies have included oxygenation of Klamath Hydroelectric Project reservoirs (MEI 2008, Horne et al. 2009, CH2M HILL 2013, Austin et al. 2016). Under the KHSA, PacifiCorp has committed to spending \$5.4 million to implement water quality improvement projects if the KHSA process reaches a critical milestone (i.e., dam removal is imminent).

## **9.5 NON-PROFIT ORGANIZATIONS**

### **9.5.1 KLAMATH WATERSHED PARTNERSHIP**

Klamath Watershed Partnership (KWP) has conducted watershed assessments for the Upper Williamson River, Upper Sprague River/Sycan River, and the Lower Sprague-Lower Williamson River, and plans to assess Upper Klamath Lake and the Klamath River. The Klamath Watershed Partnership is “a community-based non-profit organization focusing on the needs of landowners and natural resources sustainability.”<sup>4</sup> Working groups are organized in sub-watersheds and consist of community members. Through the Klamath Watershed Partnership, ranchers and landowners can get help through funding, education, and implementation of projects related to riparian restoration, grazing management, fencing, irrigation improvements, and fish screens on intake systems. KWP has been actively involved in education and outreach about stormwater and NPS pollution (KWP and ODEQ 2015). KWP also participated in the Klamath Tracking and Accounting Program (KTAP) which is an accounting system that quantifies ecosystem benefits from conservation projects through linking actions to needs of water quality improvement projects. Funding for upcoming fencing and grazing management projects will be provided by NRCS RCPP funds (see Section 9.6 below). KWP also has a beaver management program<sup>5</sup> which provides education and technical assistance to landowners for managing beaver-related issues, including onsite mitigation as well as beaver relocation.

### **9.5.2 THE NATURE CONSERVANCY**

One of The Nature Conservancy’s (TNC) primary objectives in the Klamath area is to rebuild wetlands around UKL. The Williamson Delta Wetland Preserve is TNC’s largest project in the UKL area, encompassing 5,500 acres of restored wetlands at the mouth of the Williamson River, providing rearing habitat for endangered suckers and reducing phosphorus concentrations (Wong and Hendrickson 2011). TNC has recently acquired an additional three parcels of land totaling approximately 4,000 acres along the northern shores of Agency Lake. These properties, which are referred to as the Fourmile Wetlands Preserve, are located between Fourmile Canal and Sevenmile Canal, adjacent the USFWS’s Barnes Ranch and Agency Lake Ranch properties (Hendrixson 2014). TNC has enrolled the properties in a USDA NRCS Wetland Reserves Program (WRP) easement and are re-flooding the land to restore wetlands. The Fourmile Wetlands Preserve is likely to eventually be re-connected to UKL once the dikes surrounding the Barnes Ranch and Agency Lake Ranch are breached.

### **9.5.3 TROUT UNLIMITED/KLAMATH BASIN RANGELAND TRUST**

Klamath Basin Rangeland Trust (KBRT) was founded in 2002 and merged with Trout Unlimited (TU) in 2016. Projects TU/KBRT has implemented include building wetlands on agricultural land and monitoring the effectiveness of the wetlands for nutrient removal and storage, installing

---

<sup>4</sup> <http://www.klamathpartnership.org/aboutus.html>

<sup>5</sup> <http://www.klamathpartnership.org/BMP.html>

fish screens, improving fish passage, riparian protection and enhancement, and stream reach restorations. Past projects have included assisting landowners in converting water-thirsty crops to dryland-appropriate crops and development of off-stream stock watering facilities. TU/KBRT has developed strong working relationships with landowners, NRCS, USBR, USFWS, USFS, OWEB, USGS, USBLM, UKCAN and others.

TU/KBRT is leading the development of diffuse-source treatment wetlands (see Section 11.4.1 below) in the Upper Klamath Basin (Scott 2016). Partners include landowners, USFWS, California State Coastal Conservancy, California Regional Water Boards, the Klamath Tribes, Oregon Water Resources Department, Oregon Watershed Enhancement Board (OWEB), PacifiCorp, and the National Fish and Wildlife Foundation. Two pilot projects have been implemented in the Wood River Valley and current plans are to complete approximately six more in the next few years.

## **9.6 PARTNERSHIP PROGRAMS**

### **9.6.1 KLAMATH BASIN MONITORING PROGRAM (KBMP)**

Klamath Basin Monitoring Program (KBMP) coordinates water quality monitoring in the Klamath Basin to reduce redundancy and share data, techniques, and resources (see Section 3.1 above).

### **9.6.2 UPPER KLAMATH CONSERVATION ACTION NETWORK (UKCAN)**

The Upper Klamath Conservation Action Network (UKCAN) is a partnership between the Klamath Basin Rangeland Trust (now Trout Unlimited), Klamath Soil and Water Conservation District, Klamath Watershed Partnership, Sustainable Northwest, the Klamath Tribes, The Nature Conservancy, and the Upper Klamath Water Users Association (Hendrixson and Bottcher 2015). The organizations involved in this partnership work towards habitat restoration and conservation, water use management, integrated strategic planning, project coordination, monitoring, and partnership development. The first funding source for UKCAN was a National Fish and Wildlife Foundation Keystone Initiative (Adelsberger et al. 2012). The Oregon Watershed Enhancement Board (OWEB) then approved the formation and funding of an Upper Klamath Special Investment Partnership (SIP) in 2012 (OWEB 2016). The area of focus for the SIP is from the headwaters of Upper Klamath Lake downstream to Link Dam, but also includes Spencer Creek which is a tributary to the Klamath River downstream near J.C. Boyle Reservoir.

### **9.6.3 KLAMATH REGIONAL CONSERVATION PARTNERSHIP PROGRAM (RCPP)**

Klamath Regional Conservation Partnership Program (RCPP) was approved for \$7.6 million of USDA funding in 2016 to restore wetlands and riparian habitats, improve water quality, increase instream flows in UKL tributaries, and increase farmers' drought resilience (USDA 2016b). Partners include TU/KBRT, TNC, KWP and the Klamath Lake Land Trust. The largest single project within the RCPP is the enrollment of the TNC's Fourmile Wetlands Preserve into a USDA NRCS Wetland Reserves Program (WRP) easement.

### **9.6.4 KLAMATH BASIN RESTORATION AGREEMENT (KBRA)**

The Klamath Basin Restoration Agreement (KBRA) is an agreement between Klamath Basin Tribes, irrigators, fishermen, environmental groups, counties, states, and federal agencies that

aims to both restore Klamath Basin fisheries and provide stability to the local economies. Some Consortium members signed the KBRA while others did not. The KBRA called for tens of millions of dollars for water quality improvement projects. It also called for the breaching of dikes to reconnect the Wood River Wetlands, Agency Lake Ranch, and Barnes Ranch to Agency Lake. Congress never authorized the KBRA, and thus it expired on December 31, 2015. Negotiations are ongoing, and it may or may not be replaced by a future agreement.

## 10 CONCLUSIONS

As described in this assessment, the waters within the NPS AMPP area are actually or potentially impaired from various NPS pollution sources. The NPS pollution categories in ranking of greatest concern are:

- Agriculture
- Hydromodification and Habitat Alteration

NPS pollution categories of secondary importance are:

- Forestry
- Urban

These NPS pollution categories currently contribute to water quality impairments. There are many NPS programs in place within the NPS AMPP area; however, these programs could benefit from increased coordination. NPS pollution sources will be addressed through implementation of the Clean Water Act (CWA) Section 319 NPS Management Program (see Section 11), which outlines additional short-term and long-term BMPs and program components that would be funded by various sources including, but not limited to, CWA Section 319(h) funding. Implementation of a CWA Section 319(b) NPS Management Program Plan will provide the framework for selection and implementation of best management practices and NPS pollution mitigation strategies.

## 11 NPS MANAGEMENT PROGRAM PLAN

### 11.1 MANAGEMENT PROGRAM SUMMARY

The entire NPS AMPP area is outside the reservations of the Consortium member Tribes; however, some of the area is within the ancestral territory of Shasta Indians who are enrolled members of QVIR. Consortium member Tribes have limited legal authority to mandate changes in land and water management; therefore, the Consortium's NPS AMPP relies on voluntary measures and collaboration with entities already doing work in the area.

The Consortium will work with other outside agency experts as part of its ongoing participation in multi-agency efforts on Klamath River water quality, including the Klamath Basin Monitoring Program (KBMP, see Section 3.1 above) comprised of experts from the SWRCB, NCRWQCB, USGS, EPA, USFWS, USBR, Klamath Basin Tribes and other community-based water quality monitoring groups within the Klamath Basin. The Consortium will also work with public and private landowners on the implementation of the NPS AMPP.

### 11.1.1 ADMINISTRATION

The following four paragraphs summarize the general administrative and decision-making procedures of the Consortium, which are explained in more detail in the Consortium's Strategic Plan and Bylaws (KTWQC 2015).

The Quartz Valley Indian Reservation (QVIR) is the Lead Institution within the Consortium, meaning that it is authorized to administer funds on behalf of the Consortium, and QVIR's Environmental Director is the Consortium's Chair/Program Manager. Consortium decisions are made by majority vote of the Consortium's Steering Group, which is composed of one representative of each of the Consortium's five member Tribes.

Each fiscal year, the Steering Group will develop program funding proposals to be submitted to funding entities (including but not limited to the U.S. Environmental Protection Agency) by the Chair/Program Manager on behalf of the Consortium, and allocate budgets from awarded contracts/grants into specific tasks and sub-tasks (i.e., Annual Work Plan/Budget). When mid-year adjustments are necessary, the Steering Group will determine how to re-allocate funding among tasks and sub-tasks by a majority vote of Steering Group members via email or conference call.

If it is necessary to hire consultants/contractors to implement tasks described in the Annual Work Plans/Budgets, the Lead Institution shall prepare and publish Requests for Proposals (RFPs) when appropriate. The Steering Group shall select consultants/contractors based on majority vote and the Lead Institution shall then develop contracts with the consultants/contractors.

The Steering Group shall determine the frequency of its meetings, but shall meet in person at least once yearly. The Steering Group will also conduct conference calls approximately quarterly, and communicate approximately monthly via email. Additional meetings may be called by two or more parties or at the request of the Chair/Program Manager.

The Consortium anticipates that implementation of the NPS AMPP will be part of the Consortium's Annual Work Plan. As with any other Consortium decisions, the Steering Group will decide on priorities including which projects to seek funding for in each year to implement the NPS AMPP.

Table 8 provides a summary of milestones for the management program plan.

Table 8. Management program initiation timeline and annual milestones.

<b>Activities</b>	<b>Output</b>	<b>Date</b>
Submit draft NPS Assessment and Management Program Plan to USEPA and public for review	Draft Assessment and Management Program Plan	August 5, 2016
Finalize NPS Assessment and Management Program Plan based on comments from USEPA and the public	Final Assessment and Management Program Plan	September 30, 2016
Attend spring and fall KBMP meetings to promote collaboration and coordination	2 meetings	April and November
Consortium in-person meeting to review potential projects and set priorities	1 meeting	Following start of fiscal year (typically July)
Consortium phone meetings to review potential projects and set priorities	Meeting(s)	As needed, based on grant deadlines
Develop and submit proposals to funding agencies to implement high-priority projects	Grant application	Varies according to funding source (January for EPA Tribal 319 competitive grants)

## 11.2 FUNDING SOURCES

A summary of federal and state assistance and funding resources are listed below. The program descriptions of cooperating agencies and how they relate to the abatement and control of NPS pollution within the NPS AMPP area are as follows:

### 11.2.1 FEDERAL

#### U.S. ENVIRONMENTAL PROTECTION AGENCY

The U.S. Environmental Protection Agency (USEPA) provides funding through grants administered in accordance with Section 319 of the Clean Water Act (CWA), which specifically addresses NPS pollution. These funds are available annually. There are two types of funding: base and competitive. Most Consortium member Tribes already receive base funding, which disqualifies the Consortium from also receiving base funds; however, the Consortium is eligible to apply for competitive funds. Competitive Tribal CWA 319 project implementation grants are open to proposals which are usually due in December-January. The Consortium anticipates that the EPA 319 competitive funds will be an important source for funding implementation of the NPS AMPP.

The Consortium could also potentially apply for state competitive CWA 319 funds with proposals usually due in November-December of each year.

### *U.S. DEPARTMENT OF AGRICULTURE'S NATURAL RESOURCES CONSERVATION SERVICE*

The NRCS is the primary agency responsible for providing technical, financial, and educational assistance to land users in planning and application of soil and water conservation measures. The NRCS provides funding to implement projects to minimize and prevent NPS pollution from impacting water quality. Funding is offered annually and proposals are usually due in December – January.

### *U.S. FISH AND WILDLIFE SERVICE*

The U.S. Fish and Wildlife Service (USFWS) provides competitive grants to federally recognized tribes to develop and implement programs to benefit wildlife species, including habitat restoration. The grants can be used to fund planning for wildlife and habitat conservation, fish and wildlife conservation and management actions, fish and wildlife-related laboratory and field research, natural history studies, habitat mapping, field surveys and population monitoring, habitat preservation, conservation easements, and public education that is relevant to the project. This funding is offered several times per year under different programs, often between November and April annually. Information on the USFWS Partners in Wildlife grant program for habitat restoration on private lands is available in Section 9.1.2 above.

### *NATIONAL OCEANIC AND ATMOSPHERIC ADMINISTRATION*

The National Oceanic and Atmospheric Administration (NOAA) provides for funding under the congressionally mandated Pacific Coastal Salmon Recovery Fund (PCSRF). The PCSRF funds projects, including tribal projects that improve the status of Pacific coastal salmon, prevent extinctions, and protect healthy populations. Funds are distributed annually through a competitive application process often between November and April. Until the dams are removed and salmon once again have access to the Upper Klamath Basin, PCSRF is unlikely to be a suitable source of funds for water quality restoration projects in the Upper Klamath Basin.

## 11.2.2 STATE

### *CALIFORNIA DEPARTMENT OF FISH AND WILDLIFE*

The California Department of Fish and Wildlife (CDFW) provides competitive grants under the Fisheries Restoration Grant Program for projects related to the protection/restoration of wild salmon, steelhead trout, and other fish habitats in California. Past projects funded under this program include sediment reduction projects and watershed education programs throughout coastal California. Project proposals are solicited annually, usually between October and April. As with the PCSRF, this is unlikely to be a suitable source of funds for water quality restoration projects in the Upper Klamath Basin until the dams are removed.

### *CALIFORNIA COASTAL CONSERVANCY*

The California Coastal Conservancy administers grants to fund projects that include natural resource protection and restoration in the coastal zone or affecting coastal areas. The Coastal Conservancy has funded several projects in the Upper Klamath Basin, including diffuse source treatment wetlands (Scott 2016) and a workshop to evaluate water quality improvement techniques (Stillwater Sciences et al. 2012, 2013).

### *CALIFORNIA STATE WATER RESOURCES CONTROL BOARD*

California's State Water Resources Control Board (SWRCB) allocates grant funding from the USEPA under Section 319(h) of the CWA. The funds are allocated to support implementation and planning projects that address water quality problems resulting from NPS pollution. Projects are required to be located in a watershed that has an adopted/nearly adopted Total Maximum Daily Load (TMDL) for the constituent of concern and has been identified in the NPS Program Preferences. Projects focused on working toward achieving the goals of the TMDL to restore beneficial uses will be the most competitive in the selection process.

#### OREGON WATERSHED ENHANCEMENT BOARD

The Oregon Watershed Enhancement Board (OWEB) is a state agency that provides grants to improve habitat in streams, rivers, wetlands and natural areas. OWEB has funded many habitat restoration projects in the Upper Klamath Basin. One of the larger collections of OWEB projects was the Klamath Special Investment Partnership (OWEB 2016).

#### 11.2.3 PRIVATE

As noted in Section 9.4.1, if the Klamath Hydroelectric Settlement Agreement (KHSA) proceeds as scheduled, PacifiCorp would spend up to \$5.4 million to implement Upper Klamath Basin water quality improvement projects, and up to \$560,000 per year to cover project operation and maintenance expenses related to those projects. The list of projects would be developed by the Interim Measures Implementation Committee (IMIC) and approved by the Oregon Department of Environmental Quality (ODEQ) and California's State and Regional Water Boards.

### **11.3 CATEGORIES OF NONPOINT SOURCE POLLUTION**

In the NPS assessment (see sections 1 through 10 above), the Consortium identified categories of NPS pollution that are likely impacting water quality, as shown below. The Consortium then ranked the identified categories of NPS pollution based on their relative importance.

The following two categories are high priority, due to the widespread extent and severity of impacts:

- Agriculture (AGR)
- Hydromodification and Habitat Alteration (HHA)

The following two categories are much lower priority given their lesser contribution to NPS pollution within the NPS AMPP area:

- Forestry (FOR)
- Urban (URB)

The Consortium's strategy focuses on preventing NPS pollution and treating the sources of water quality impairment; however, under a category entitled Other (OTH), we also include tasks and BMPs to strategically target the symptoms of excessive eutrophication which result from multiple sources.



#### **11.4 TASKS AND BMPS**

In order to implement an effective, comprehensive NPS management program, all sources of pollution must be addressed in a manner that provides alternatives and flexibility. This document is designed to present a proposed strategy with flexibility in management and implementation. The goals and tasks outlined can be utilized, altered, and, prioritized to protect and restore water quality.

The tasks and BMPS listed in the following section are the Consortium's recommendations for improving water quality within the Upper Klamath Basin. The BMPS are categorized as either short-term or long-term tasks. Short-term tasks are defined as tasks that can be implemented immediately and implementation consists of defined tasks that will facilitate control on NPS pollution. Long-term tasks are defined as tasks that either require precursor tasks to be completed before implementation, or are ongoing management or operational changes.

Implementation of the following schedule (Figure 10) depends on the availability of adequate funding pursuant to Section 319 of the CWA and other applicable sources. The scope of the schedule is intended to err on the side of being comprehensive and it is unlikely that sufficient funds will be available to implement all tasks on the schedule. The Consortium reserves the right to alter or modify the schedule based on immediate needs, resources available, and economics. Priority will be given to projects that reduce phosphorus concentrations into Upper Klamath Lake, with the ultimate goal of restoring water quality within UKL and the Klamath River downstream.

Table 9. NPS implementation schedule by fiscal year (July 1 – June 30). Tasks are sorted first by NPS category and then by goal. Key to NPS categories: Agriculture (AGR), Hydromodification and Habitat Alteration (HHA), Forestry (FOR), Urban (URB), and Other (OTH).

Goal	Task	Year 1 (2016/ 2017)	Year 2 (2017/ 2018)	Year 3 (2018/ 2019)	Year 4 (2019/ 2020)	Year 5 (2020/ 2021)
Goal: Reduce runoff of phosphorus from land due to agricultural activities and prevent phosphorus delivery to important aquatic habitats downstream	Short-Term Task AGR-1: Restore riparian corridors to minimize transport of sediment and phosphorus into waterbodies	x	x	x	x	x
	Short-Term Task AGR-2: Implement pilot-scale diffuse-source treatment wetlands and monitor effectiveness	x	x	x		
	Long-Term Task AGR-3: Implement many diffuse-source treatment wetlands	x	x	x	x	x
	Short-Term Task AGR-4: Find suitable locations and develop designs for large-scale treatment wetlands	x	x	x	x	x
	Long-Term Task AGR-5: Implement large-scale treatment wetlands	x	x	x	x	x
	Short-Term Task AGR-6: Evaluate effects and develop designs for reuse and recirculation of agricultural drain water	x	x	x	x	x
	Long-Term Task AGR-7: Implement projects to reuse and recirculate agricultural drain water	x	x	x	x	x
	Long-Term Task AGR-8: Implement projects to improve tailwater management	x	x	x	x	x
	Long-Term Task AGR-9: Education and technical assistance to farmers and ranchers	x	x	x	x	x
Goal: Increase instream flows by reducing irrigation demand	Long-Term Task AGR-10: Convert irrigated pastures to dryland agriculture	x	x	x	x	x
	Long-Term Task AGR-11: Reduce summer irrigation and improve grazing management	x	x	x	x	x
Goal: Restore riparian corridors and promote floodplain connectivity	Short-Term Task HHA-1: Develop parcel-specific riparian restoration plans	x	x			
	Long-Term Task HHA-2: Implement projects to restore riparian areas by constructing riparian fencing, planting riparian vegetation, and providing livestock with off-channel water sources	x	x	x	x	x
	Long-Term Task HHA-3: Mechanical manipulation to remove levees and restore channel sinuosity	x	x	x	x	x
	Long-Term Task HHA-4: Implement a beaver management program and consider deployment of beaver dam analogues	x	x	x	x	x

Goal	Task	Year 1 (2016/ 2017)	Year 2 (2017/ 2018)	Year 3 (2018/ 2019)	Year 4 (2019/ 2020)	Year 5 (2020/ 2021)
Goal: Restore wetlands around UKL and tributaries to improve water quality and habitat	Short-Term Task HHA-5: Develop wetland restoration plans	x	x	x	x	x
	Long-Term Task HHA-6: Implement wetland restoration plans	x	x	x	x	x
Goal: Eliminate effects of Klamath Hydroelectric Project on water temperature, algal toxins, and food web	Short-Term Task HHA-7: Develop plan and permitting to remove J.C. Boyle, Copco 1/2, and Iron Gate dams	x	x	x	x	x
	Long-Term Task HHA-8: Implement plan to remove J.C. Boyle, Copco 1/2, and Iron Gate dams				x	x
Goal: Reduce impacts of private forest management on aquatic habitats	Long-Term Task FOR-1: Improve riparian protections in Oregon Forest Practice Rules	x	x	x	x	x
Goal: Reduce sediment runoff from forest roads	Short-Term Task FOR-2: Conduct road assessments	x	x	x	x	x
	Long-Term Task FOR-3: Upgrade and decommission roads	x	x	x	x	x
Goal: Reduce delivery of urban stormwater contaminants to important aquatic habitats downstream	Long-Term Task URB-1: Provide outreach, education, and technical assistance to residents and businesses in Klamath Falls	x	x	x	x	x
	Long-Term Task URB-2: Minimize effects of new development on stormwater	x	x	x	x	x
Goal: Reduce organic matter loads discharged from UKL into Klamath River	Short-Term Task OTH-1: Develop conceptual layout for system to remove algal biomass at outlet of UKL	x	x			
	Short-Term Task OTH-2: Conduct pilot-scale test to remove algal biomass at outlet of UKL	x	x			
	Long-Term Task OTH-3: Develop detailed design and engineering for system to remove algal biomass at outlet of UKL and obtain necessary permits		x	x		
	Long-Term Task OTH-4: Construct/operate full-scale system to remove algal biomass at outlet of UKL		x	x	x	x

#### 11.4.1 AGRICULTURE

*GOAL: REDUCE RUNOFF OF PHOSPHORUS FROM LAND DUE TO AGRICULTURAL ACTIVITIES AND PREVENT PHOSPHORUS DELIVERY TO IMPORTANT AQUATIC HABITATS DOWNSTREAM*

*Long-Term Task AGR-1: Restore riparian corridors to minimize transport of sediment and phosphorus into waterbodies*

Many of the rivers and streams upstream of UKL have poor riparian function, resulting in erosion and delivery of phosphorus into waterbodies. Transforming these degraded reaches into robust riparian corridors is essential to reducing the inflow of phosphorus into UKL as well as generally improving aquatic habitats in the tributaries. Tasks and BMPs for restoring riparian function are listed below in the Section titled “11.4.2.”

*Short-Term Task AGR-2: Implement pilot-scale diffuse-source treatment wetlands and monitor effectiveness*

Diffuse-Source Treatment Wetlands (DSTWs) consist of multiple small-scale constructed wetlands placed throughout a given catchment rather than one large treatment wetland at the bottom of the catchment (Stillwater Sciences et. al. 2012, 2013). DSTWs are intended to provide on-site removal of sediment, nutrients, and herbicides/pesticides, particularly for first-flush runoff events. Ancillary benefits may include a reduction in peak flows.

DSTWs are essentially small-scale wetlands scattered throughout the watershed, and thus the basic design elements are similar to those of larger treatment wetlands. DSTWs in pastures and agricultural fields can be operated either as continuous flow-through systems or as intermittent flow-through systems. Given the importance of downstream water use in the Klamath Basin, these systems would not be designed as terminal wetlands, but rather they would treat on-site runoff such that there would be an outflow of water from each site. Since DSTWs are placed within agricultural fields, exclusion fencing is required to prevent cattle damage. Ideal locations of diffuse-source wetlands would be in areas with remnant wetlands that possess hydric soils and are in proximity to irrigation ditches and drains. Additional information on DSTWs, including cost estimates, can be found in Stillwater Sciences et al. (2012 and 2013).

As noted in Section 9.5.3 above, Trout Unlimited and its partners are developing DSTWs in the Upper Klamath Basin, beginning with a few intensively monitored pilot projects in the Wood River Valley (Scott 2016). These pilot projects will provide data on effectiveness and hopefully lead to improved designs. There is also a need to evaluate the performance of DSTWs in other locations such as the Sprague River Valley which is the other primary agricultural area upstream of Upper Klamath Lake.

*Long-Term Task AGR-3: Implement many diffuse-source treatment wetlands*

If the initial demonstration projects in the Wood and Sprague valleys indicate that DSTWs are effective in reducing phosphorus at a relatively low per-unit cost, the Consortium would like to support the construction of many DSTW's throughout the Wood and Sprague valleys.

*Short-Term Task AGR-4: Find suitable locations and develop designs for large-scale treatment wetlands*

The larger the area of a treatment wetland, the greater the incoming amounts of nutrients and water the wetland can effectively treat (CH2M HILL 2012). Given the large volumes of water in the Klamath Basin, very large wetlands (on the order of thousands of acres or tens of thousands of acres) would be necessary to have basin-scale effects, but smaller-scale wetlands could have local effects. PacifiCorp and its consultants have conducted several treatment wetland evaluations including evaluating potential locations where wetlands could be sited (Lyon et al. 2009, CH2M HILL 2012) and a conceptual design for a demonstration wetland facility which could be used to quantify nutrient removal rates and water balances (CH2M HILL 2014). The USGS and its collaborators have modeled the potential water quality effects of adding treatment wetlands along Keno Reservoir (Sullivan et al. 2014).

The deployment of large-scale treatment wetlands in the Upper Klamath Basin is currently hampered by water rights (and water availability) issues (Stillwater Sciences et. al. 2012, 2013) and a lack of willing landowners; however, the Consortium still favors consideration of such wetlands given their potential to improve water quality. From the perspective of reducing phosphorus loading to UKL, areas adjacent to the lower reaches of Fourmile Canal, Sevenmile Canal, the Wood River, and Williamson River would be excellent locations for treatment wetlands, but in the Klamath Basin Restoration Agreement (KBRA) much of that area was slated for levee breaching and reconnection to UKL which limits the ability to use that area for highly managed treatment wetlands. USBR (2016b) evaluated the potential for using treatment wetlands to reduce nutrient loads in the Klamath Straits Drain prior to discharge into Keno Reservoir. Other potential locations to consider for treatment wetlands include areas adjacent to Keno Reservoir such as the Miller Island Wildlife Refuge.

*Long-Term Task AGR-5: Implement large-scale treatment wetlands*

If water rights, water availability, and landowner willingness issues can be addressed, suitable areas identified, and designs developed, then treatment wetlands should be implemented at a large scale.

*Short-Term Task AGR-6: Evaluate effects and develop designs for reuse and recirculation of agricultural drain water*

Agricultural drains such as the Klamath Straits Drain can have very high nutrient concentrations (Stillwater Sciences et. al. 2012). The discharge of such water into the Klamath River can increase nutrients concentrations in the river. One alternative to discharge of drain water into the river is to reuse it by recirculating it back into irrigation canals. USBR (2016b) identified several potential areas where recirculation may be feasible. Due to evaporative concentration of salts in soils, salinity is a potential issue that would need to be taken into consideration in any recirculation project. One alternative is to divert drain water into a reservoir during the summer where it can be held for months and then later discharged to the river during later fall or winter when water quality is less impaired and nutrients will have less effect (i.e., due to cold temperatures and low solar energy, algal growth is limited during the winter regardless of nutrient conditions). The Consortium favors the evaluation of reuse and recirculation projects including development of project designs and engineering.

*Long-Term Task AGR-7: Implement projects to reuse and recirculate agricultural drain water*

If reuse and recirculation projects can be identified and designed to benefit water quality without negatively affecting instream flows, the Consortium would consider supporting implementation of such projects.

*Long-Term Task AGR-8: Implement projects to improve tailwater management*

Tailwater is water that runs off the lower end of a field during flood (surface) irrigation (Schwankl et al. 2007). Production of some tailwater is a normal and nearly unavoidable part of flood irrigation, but can detrimentally affect water quality if the water heats up and transports organic matter and nutrients into downstream waterbodies (Aqua Terra Consulting 2011). Flood irrigation is a common practice in Upper Klamath Basin pastures (USDA 2009) and improved tailwater management has been recommended as a component of efforts to reduce phosphorus loading to Upper Klamath Lake (Walker et al. 2015). Methods for minimizing the effects of tailwater on water quality include careful management, scheduling, capture and reuse (SVRCD no date, Schwankl et al. 2007). Tailwater can also be treated using Diffuse Source Treatment Wetlands (DSTWs).

The Consortium generally supports implementation of projects to minimize the implementation of projects to improve tailwater management, particularly upstream of Upper Klamath Lake; however, it is important to consider to net effects of tailwater projects on watershed-scale water balances. For example, a project to capture tailwater (which currently flows back to a stream via overland flow and subsurface infiltration) and use it to irrigate a pasture that is not currently irrigated would likely result in reduced instream flows due to increased evapotranspiration (i.e., consumptive use) in the newly irrigated field.

*Long-Term Task AGR-9: Education and technical assistance to farmers and ranchers*

Education and technical assistance are critically important to improving water quality in the Upper Klamath Basin. The Consortium encourages educational efforts to reduce detrimental effects of land and water management on water quality and habitat. Agencies involved with education and outreach to farmers and ranchers include the Natural Resources Conservation Service (NRCS), Klamath Soil and Water Conservation District (KSWCD), Oregon Department of Agriculture (ODA), Oregon Department of Environmental Quality (ODEQ), Klamath Basin Research & Extension Center (KBREC), Klamath Watershed Partnership (KWP), and Trout Unlimited (TU).

**GOAL: INCREASE INSTREAM FLOWS BY REDUCING IRRIGATION DEMAND**

*Long-Term Task AGR-10: Convert irrigated pastures to dryland agriculture*

The Upper Klamath Basin Comprehensive Agreement (UKBCA, see Section 9.2.1 above) calls for inflows to Upper Klamath Lake to be increased by 30,000 acre-feet per year to be achieved by reducing the net consumptive use of water for irrigated agriculture (UKBCA 2014). An important mechanism for reducing the net consumptive use is to reduce the irrigated area. The Natural Resources Conservation Service, Klamath Soil and Water Conservation District, and local ranchers have developed and tested a seed mix and multi-year process to establish productive dryland pasture which does not require ongoing irrigation beyond the initial establishment period (Ferguson and Watkins 2015, OWEB 2016, USDA 2016a). In addition to

conserving water quantity, dryland pasture should also improve water quality by reducing tailwater. The Consortium favors the conversion of irrigated pastures to dryland agriculture.

*Long-Term Task AGR-11: Reduce summer irrigation and improve grazing management*

An alternative to reducing irrigated area is to apply less water to irrigated lands, such as by reducing the number of flood irrigation applications per year (KBRT 2011). Experiments in the Wood River valley showed that reduced irrigation (one irrigation event per summer) and improved grazing management (rotating cattle between pastures with a 30-day rest period between grazing cycles) only reduced forage production by 5% compared to conventional management (NRCS 2010, KBRT 2011). The Consortium supports reduced irrigation and improved grazing management as a method to reduce irrigation demand.

#### 11.4.2 HYDROMODIFICATION AND HABITAT ALTERATION

*GOAL: RESTORE RIPARIAN CORRIDORS AND PROMOTE FLOODPLAIN CONNECTIVITY*

*Short-Term Task HHA-1: Develop parcel-specific riparian restoration plans*

The Upper Klamath Basin Comprehensive Agreement (UKBCA, see Section 9.2.1 above) calls for the establishment of Riparian Management Areas along streams upstream of Upper Klamath Lake and a Riparian Action Plan including parcel-specific management and restoration actions which will then be incorporated into a Riparian Management Agreement for each property within the Riparian Management Corridor (UKBCA 2014). The Consortium favors the approach outlined in the UKBCA's riparian program.

*Long-Term Task HHA-2: Implement projects to restore riparian areas by constructing riparian fencing, managing livestock access, planting riparian vegetation, and providing livestock with off-channel water sources*

Once parcel-specific restoration plans are developed, they should be implemented. It is expected that management measures will include construction of riparian fencing, managing livestock access to riparian areas, providing livestock with off-channel sources of water and salt, and planting to encourage establishment of riparian vegetation. These efforts should be informed and guided by previous evaluations of restoration projects in the Upper Klamath Basin, including the NewFields and Kondolf (2012) report.

*Long-Term Task HHA-3: Mechanical manipulation to remove levees and restore channel sinuosity*

In some cases, such as where streams are strongly confined by armored levees and/or incised such as the South Fork of the Sprague River, natural geomorphic processes are sufficiently altered that passive restoration is unlikely to be successful or will take a very long time (O'Connor et al. 2015). In such cases, mechanical manipulations such as levee setbacks, excavation of new channels, or floodplain grading, may be necessary. It is expected that these areas will be identified in the parcel-specific restoration plans. Restoring floodplain connectivity in these areas will promote deposition of sediment onto floodplains and reduce downstream transport of sediment-bound phosphorus. Mechanical manipulations can be extremely costly, so they should only be used when they are the only option likely to be effective. The Consortium recommends that beaver dam analogues (discussed in the following section) be evaluated as a low-cost technique for reversing stream channel incision by aggrading stream channels.

*Long-Term Task HHA-4: Implement a beaver management program and consider deployment of beaver dam analogues*

American beaver (*Castor canadensis*) are a keystone species which can profoundly alter stream ecosystems (Pollock et al. 2015). They are native to the Klamath Basin and are present throughout the basin (Friedrichsen 1996, Silloway 2010, Lanman et al. 2013). Dams created by beavers can retain sediment, reverse channel incision, restore floodplain connectivity, promote habitat complexity, increase channel sinuosity, create thermal diversity, recharge groundwater, and raise groundwater elevation (Pollock et al. 2014, Majerova et al. 2015). As a result, beavers and their dams can greatly enhance habitat quality for anadromous salmonids, including coho salmon (Pollock et al. 2004) and steelhead (Bouwes et al. 2016). An intensively monitored seven-year experiment in Oregon demonstrated significant increases in the density, survival, and production of juvenile steelhead following installation of beaver dam analogues (BDAs) which mimic the functions of natural beaver dams (Bouwes et al. 2016).

Beavers can cause problems for humans in floodplain areas by consuming crops and landscaping, flooding roads and property, and damming irrigation infrastructure. Non-lethal and effective methods have been developed for addressing many of these issues (Pollock et al. 2015). Recognizing the water supply and ecosystem benefits beavers provide, the Klamath Watershed Partnership has been providing education and technical assistance to landowners for managing beaver issues, including on-site mitigation as well as beaver relocation (though only a few relocations have been conducted due to stringent state regulations). The Consortium supports efforts to maintain and expand beaver populations, and considers beaver dam analogs to be a promising cost-effective technique for stream restoration.

*GOAL: RESTORE WETLANDS AROUND UKL AND TRIBUTARIES TO IMPROVE WATER QUALITY AND HABITAT*

*Short-Term Task HHA-5: Develop wetland restoration plans*

Most of the historical wetlands around Upper Klamath Lake (UKL) were diked, drained, and converted to agriculture (see Section 5.2 above). Some of those historic wetlands, such as the mouths of the Wood and Williamson rivers, are now being restored back to wetlands; however, restoration of additional areas is necessary to help improve habitat and water quality conditions in UKL and the Klamath River downstream. Entities involved in restoring wetlands around UKL include The Nature Conservancy (TNC), U.S. Bureau of Land Management, U.S. Fish and Wildlife Service, the U.S. Bureau of Reclamation, Trout Unlimited, and the Klamath Tribes. The Klamath Basin Restoration Agreement (KBRA, see Section 9.6.4) called for the breaching of dikes to reconnect the Wood River Wetlands, Agency Lake Ranch, and Barnes Ranch to Agency Lake, with the goal of increasing water storage in UKL and restoring wetland habitat. TNC recently acquired approximately 4,000 acres adjacent to Agency Lake Ranch and Barnes Ranch which it is in the process of restoring back to wetlands, supported in part by the USDA NRCS Wetland Reserves Program (see Section 9.5.2 and 9.6.3). The Consortium encourages those entities involved in wetland restoration to continue their work, and to expand the effort if additional areas become available. An essential step is the development of wetland restoration plans and designs for specific parcels.

*Long-Term Task HHA-6: Implement wetland restoration plans*

Once parcel-specific wetland restoration plans are developed, they should be implemented.



GOAL: ELIMINATE EFFECTS OF KLAMATH HYDROELECTRIC PROJECT ON WATER TEMPERATURE, ALGAL TOXINS, AND FOOD WEB

*Short-Term Task HHA-7: Develop plan and permitting to remove J.C. Boyle, Copco 1/2, and Iron Gate dams*

The reservoirs in the Klamath Hydroelectric Project (KHP) contribute to water quality impairments in the Klamath River (see Section 0 above). The Klamath Hydroelectric Settlement Agreement (KHTSA) is a multi-party agreement to remove J.C. Boyle, Copco 1, Copco 2, and Iron Gate dams. All Consortium member Tribes favor removing these dams, but only some Consortium member Tribes support the KHTSA as the vehicle for achieving that goal. Regardless of what path is ultimately chosen for dam removal, the process will be complicated and take years of planning and studies including development of required environmental permits.

*Long-Term Task HHA-8: Implement plan to remove J.C. Boyle, Copco 1/2, and Iron Gate dams*

Once plans are developed for how to remove the dams and appropriate permits are obtained, the dams should be removed. The KHTSA targets 2020 as the year in which dam removal would occur.

#### 11.4.3 FORESTRY

Relative to agriculture (Section 11.4.1) and hydromodification and habitat alteration (Section 11.4.2), forestry is a small contributor to basin-scale NPS pollution in the Upper Klamath Basin but for the sake of completeness we include tasks to address it.

GOAL: REDUCE IMPACTS OF FOREST MANAGEMENT ON AQUATIC HABITATS

*Long-Term Task FOR-1: Improve riparian protections in Oregon Forest Practice Rules*

Riparian forests are important to streams because they stabilize streambanks, provide shade, and contribute wood which provides habitat structure. Anadromous fish are currently blocked from accessing the Upper Klamath Basin by a series of dams on the mainstem Klamath River (see Section 4.3 above). Given that these dams are likely to be removed within the next decade through the KHTSA or other means, it is important to start protecting and restoring habitat within the tributaries upstream of Iron Gate Dam. One tributary of particular importance is Spencer Creek, which has very high intrinsic potential to serve as coho salmon habitat but is currently a tributary to J.C. Boyle Reservoir (NMFS 2014). If streamflow and riparian conditions in Spencer Creek were restored to natural conditions then water temperatures would be suitable for coho salmon rearing throughout almost the entire length of the creek (ODEQ 2010a). One factor that may retard recovery of aquatic habitats in these tributaries is weak regulation of private timberland in Oregon. Reviews of the Oregon Forest Practice Rules have found that they do not adequately protect riparian areas (NMFS 2014). Protections for small and medium-sized streams are particularly deficient (IMST 1999). Following experiments showing that harvest under the current rules increased stream temperatures, Oregon regulators initiated a process to increase riparian protections; however, the Klamath Basin is not included in the proposed revisions. Due to the Northwest Forest Plan and the Aquatic Conservation Strategy (USFS and BLM 1994), riparian protections on federal lands are much stronger than those on private lands in Oregon. The Consortium has no legal authority to alter the Oregon Forest Practice Rules; however, it can recommend changes when opportunities for public comment arise. If the Klamath Basin Restoration Agreement (KBRA, see Section 9.6.4 above) is adopted, Oregon Department of

Forestry's ability to strengthen water quality regulatory requirements on private forestry operations within the Klamath Basin may be limited<sup>6</sup>.

*Short-Term Task FOR-2: Conduct road assessments*

Some areas within the Upper Klamath Basin have dense networks of roads which can produce sediment and impair streams. Field surveys of roads are necessary to determine which sections of roads have the potential to cause the most problems. Priority areas for road assessments would be the watersheds of tributaries with the highest potential to support anadromous salmonids.

*Long-Term Task FOR-3: Upgrade and decommission roads*

The highest-priority sites identified within the assessed areas would be targeted for upgrade treatments to reduce their potential to generate sediment. Treatments may include out sloping and culvert replacement. Roads that are no longer necessary would be considered for decommissioning (i.e., removal).

#### 11.4.4 URBAN

Given the limited urbanized area, urban stormwater is a minor contributor to NPS pollution within the Upper Klamath Basin and its impacts to water quality are far smaller than agriculture (Section 11.4.1) and hydromodification and habitat alteration (Section 11.4.2), but for the sake of completeness we include tasks to address it.

*GOAL: REDUCE DELIVERY OF URBAN STORMWATER CONTAMINANTS TO IMPORTANT AQUATIC HABITATS DOWNSTREAM*

*Long-Term Task URB-1: Provide outreach, education, and technical assistance to residents and businesses in Klamath Falls*

Despite the fact that urban stormwater contributes only a small fraction of the NPS pollution within the Upper Klamath Basin, educating urban residents about stormwater is important. Stormwater education contributes to residents' understanding of their role in the watershed and promotes appreciation and connection to rivers, lakes, and wildlife habitat. The Klamath Watershed Partnership has been actively involved in education and outreach about stormwater and NPS pollution (KWP and ODEQ 2015). The Consortium will seek to partner with KWP and other organizations promoting education and outreach about NPS pollution.

*Long-Term Task URB-2: Minimize effects of new development on stormwater*

Given the small contribution of urban stormwater to NPS pollution in the Upper Klamath Basin, the Consortium does not recommend costly retrofits to existing urban infrastructure to reduce NPS pollution; however, the Consortium does recommend that future urban development include Lower Impact Development (LID)<sup>7</sup> principles to increase infiltration and decrease stormwater

---

<sup>6</sup> Excerpt from the KBRA "25.1.3. Forestry: Private forestry operations complying with water protection rules administered by the Oregon Department of Forestry, and with rule amendments, if any, adopted to implement the Fisheries Program, shall not be subject to further water quality requirements under Oregon Revised Statutes chapter 468B or 527, if any, arising solely from reintroduction and the designation or presence of new fish beneficial uses.

<sup>7</sup> LID is more commonly referred to as "Low Impact Development," but development conducted with LID principles still has impacts, and those impacts can in fact be great depending on the scale and location. As noted by

runoff. LID features can be included in new development at relatively low cost if incorporated early in the design process. The only mention of stormwater runoff in Klamath County's current land development code<sup>8</sup> is a single sentence that states "The purposes of landscaping requirements are to maintain and enhance the appearance of structures and properties, to provide for visual privacy and a quality visual environment, and to provide areas on sites to absorb rainfall and reduce stormwater runoff." The land development code lacks the key LID principle that impervious surfaces (e.g., roofs and parking lots) should be sloped such that water is routed to landscaped swales and depressions where it can infiltrate. To minimize stormwater runoff, landscaped areas should not be islands hydrologically disconnected (i.e., elevated above the impervious area and separated by curbs) within the developed area, and impervious areas should not be sloped to concentrate water into hardened drains which rapidly carry water offsite.

#### 11.4.5 OTHER

Tasks in this section are interventions to address the symptoms of excessive algal productivity in Upper Klamath Lake from Keno Reservoir downstream. These tasks are assigned to the "other" category because these symptoms are a result of multiple interacting categories of NPS pollution.

#### *GOAL: REDUCE ORGANIC MATTER LOADS DISCHARGED FROM UKL INTO KLAMATH RIVER*

##### *Short-Term Task OTH-1: Develop conceptual layout for system to remove algal biomass at outlet of UKL*

Several previous projects have evaluated the potential for removal of algal biomass near Link Dam to improve water quality in Keno Reservoir downstream. Technologies evaluated thus far included stormwater technology (hydrodynamic separators, Watercourse Engineering, Inc. 2013, 2014a, 2014b) and algal harvesting technologies (Stillwater et al. 2012, 2013; PacifiCorp 2015; Carlson and Hughes 2016). Removing algae would be beneficial for several other reasons: 1) modest direct reduction in downstream nutrients because algae contains phosphorus (Sullivan et al. [2013] predict that a 50% reduction in algae and particulate organic matter would reduce June-October Keno Reservoir total phosphorus loads by ~8%), 2) potential for creation of useful products (e.g., pharmaceuticals, human/animal dietary supplements, or fertilizer) if it is cost-effective to do so, and 3) in contrast to a chemical treatment like alum, algal harvesting does not produce potentially toxic byproducts.

Carlson and Hughes (2016) coordinated a series of conference calls with the IMIC (see Section 9.4.1 above) to discuss key questions about algal harvesting and summarized the resulting discussions. Participants generally agreed that removing 25% of the algae at Link Dam was likely the approximate upper limit of feasibility. The Sullivan et al. (2013) model scenarios predict that a 25% reduction in algae and organic matter would increase average June 15-October 31 dissolved oxygen conditions in Keno Reservoir by approximately 0.8 mg/L. This increase would notably improve water quality, but is far short of meeting dissolved oxygen criteria. In summary, algae removal has the potential to make some limited improvement in Keno Reservoir dissolved oxygen, but cannot by itself resolve the oxygen depletion problem.

---

Strecker (2001), "Low Impact Development" is a misnomer and therefore instead we prefer the term "Lower Impact Development."

<sup>8</sup> [http://www.klamathcounty.org/depts/planning/downloads/Codes\\_Plans/LandDevCode/LDCChapter60.pdf](http://www.klamathcounty.org/depts/planning/downloads/Codes_Plans/LandDevCode/LDCChapter60.pdf)

The Consortium recommends that algae removal be further evaluated. The logical next step would be to evaluate conceptual layouts and develop rough estimates of effectiveness and costs. The two largest obstacles to implementing algae harvesting are environmental permitting (due to presence of endangered fish) and cost (due to the large scales required). One of the largest potential costs would be the construction of a fish screen to minimize take of endangered fish. There may be ways to minimize costs to take advantage of existing infrastructure such as the A-Canal fish screen or the Eastside Forebay at Link Dam.

*Short-Term Task OTH-2: Conduct pilot-scale test to remove algal biomass at outlet of UKL*

Pilot tests are necessary to assess the potential harvested amounts and per-unit costs for algal harvesting systems. Previous efforts (Stillwater et al. 2012, 2013) already tried to get this information from the literature and there is not much information available. In-lake pilot tests are absolutely critical. Ideally, several different technologies should be tested. Due to permitting requirements, the Consortium envisions the pilot tests as a two-year project with the first year being laying the groundwork for a pilot test that would be conducted in the second year. That groundwork should include recruiting algal harvesters to participate, creating a study design, and obtaining necessary permits.

*Long-Term Task OTH-3: Develop detailed design and engineering for system to remove algal biomass at outlet of UKL and obtain necessary permits*

If the pilot projects and conceptual designs indicate that algae harvesting would be feasible and cost-effective, then a full-scale system should be designed and engineered and permits acquired.

*Long-Term Task OTH-4: Construct/operate full-scale system to remove algal biomass at outlet of UKL*

Once the full-scale algal harvesting system is designed and permits have been acquired, then the system should be constructed and put into operation.

## 12 REFERENCES

Adelsberger, C., N. Athearn, M. Barry, US Fish and Wildlife Service, J. Bottcher, K. Fischer, H. Hendrixson, C. Lambert, G. Monroe, K. Ryals, and D. Watson. 2012. Business Plan for the Upper Klamath Basin Keystone Initiative, a 10-Year Initiative to Secure Upper Klamath Basin Native Fish Populations: Lost River Sucker, Shortnose Sucker, and Klamath Redband Rainbow Trout. Report prepared for the National Fish and Wildlife Foundation, Washington, D. C.

Aqua Terra Consulting, 2011. Shasta River Tailwater Reduction Plan. Prepared for the Shasta Valley Resource Conservation District by Aqua Terra Consulting, Mt. Shasta, CA. 32 p. [https://www.dropbox.com/s/onsgjdavlb8h2y5/Tailwater-Reduction-Plan\\_9\\_11.pdf](https://www.dropbox.com/s/onsgjdavlb8h2y5/Tailwater-Reduction-Plan_9_11.pdf)

- Asarian, E. and J. Kann. 2013. Synthesis of Continuous Water Quality Data for the Lower and Middle Klamath River, 2001-2011. Prepared by Kier Associates and Aquatic Ecosystem Sciences for the Klamath Basin Tribal Water Quality Work Group. 50p. + appendices. [http://www.riverbendsci.com/reports-and-publications-1/Klamath\\_2001\\_2011\\_sonde\\_rpt\\_20130502\\_final.pdf](http://www.riverbendsci.com/reports-and-publications-1/Klamath_2001_2011_sonde_rpt_20130502_final.pdf)
- Asarian, E., J. Kann, and W. Walker. 2009. Multi-year nutrient budget dynamics for Iron Gate and Copco Reservoirs, California. Prepared by Riverbend Sciences, Kier Associates, Aquatic Ecosystem Sciences, and William Walker for Karuk Tribe Department of Natural Resources, Orleans, California.
- Asarian, J.E. and J.D. Walker. 2016. Long-Term Trends in Streamflow and Precipitation in Northwest California and Southwest Oregon, 1953-2012. *JAWRA Journal of the American Water Resources Association* 52:241–261. doi: 10.1111/1752-1688.12381.
- Austin, D., M. Deas, and K. Carlson. 2016. (Draft) Technical Memorandum: Interim Measure 11, Activity 4 – Conceptual Feasibility Study of Oxygenation Systems at Keno Reservoir, April 7, 2016. Prepared by CH2M and Watercourse Engineering for PacifiCorp, Portland, Oregon.
- Barbiero, R. P., and J. Kann. 1994. The importance of benthic recruitment to the population development of *Aphanizomenon flos-aquae* and internal loading in a shallow lake. *Journal of Plankton Research* 16: 1581–1588.
- Bouwes, N., N. Weber, C.E. Jordan, W.C. Saunders, I.A. Tattam, C. Volk, J.M. Wheaton, and M.M. Pollock. 2016. Ecosystem Experiment Reveals Benefits of Natural and Simulated Beaver Dams to a Threatened Population of Steelhead (*Oncorhynchus mykiss*). *Scientific Reports* 6:28581. doi: 10.1038/srep28581.
- Bradbury, J.P., Colman, S.M. and R.L. Reynolds. 2004. The history of recent limnological changes and human impact on Upper Klamath Lake, Oregon. *Journal of Paleolimnology* 31: 151-161.
- Carlson, K., and B. Hughes. 2016. Technical Memorandum: Interim Measure 11, Activity 7 – Assessment of Potential Algae Harvesting and Removal Techniques at Link River Dam, July 12, 2016. Prepared by CH2M for PacifiCorp, Portland, Oregon.
- Carpenter, K.D., D.T. Snyder, J.H. Duff, F.J. Triska, K.K. Lee, R.J. Avanzino, and S. Sobieszczyk. 2009. Hydrologic and water-quality conditions during restoration of the Wood River Wetland, Upper Klamath River Basin, Oregon, 2003-2005. *Scientific Investigations Report 2009-5004*. U.S. Geologic Survey.
- Ciotti, D., S.M. Griffith, J. Kann, and J. Baham. 2010. Nutrient and sediment transport on Flood-Irrigated pasture in the Klamath Basin, Oregon. *Rangeland Ecology and Management* V 63 (3): 308-316.
- CH2M HILL. 2012. Approaches to Water Quality Treatment by Wetlands in the Upper Klamath Basin. Prepared for PacifiCorp Energy, Portland, OR. Prepared by CH2M HILL, Inc., Portland, OR. August 2012. [http://www.pacificorp.com/content/dam/pacificorp/doc/Energy\\_Sources/Hydro/Hydro\\_Licensing/Klamath\\_River/2012Aug\\_WQTreatmentWetlandsFinal.pdf](http://www.pacificorp.com/content/dam/pacificorp/doc/Energy_Sources/Hydro/Hydro_Licensing/Klamath_River/2012Aug_WQTreatmentWetlandsFinal.pdf)

- CH2M HILL. 2013. Assessment of Technologies for Dissolved Oxygen Improvement in J.C. Boyle Reservoir. Final Report. Prepared for PacifiCorp Energy, Portland, Oregon. Prepared by CH2M HILL, Inc. July 2013.  
[http://www.pacificorp.com/content/dam/pacificorp/doc/Energy\\_Sources/Hydro/Hydro\\_Licensing/Klamath\\_River/JCB%20DO%20Assessment%20Final%20Report%20\(July%202013\).pdf](http://www.pacificorp.com/content/dam/pacificorp/doc/Energy_Sources/Hydro/Hydro_Licensing/Klamath_River/JCB%20DO%20Assessment%20Final%20Report%20(July%202013).pdf)
- CH2M HILL. 2014. Demonstration Wetland Facility Preliminary Research and Implementation Plan Klamath River, Oregon. 2014. Prepared for PacifiCorp by CH2M HILL, Portland, OR.  
[http://www.pacificorp.com/content/dam/pacificorp/doc/Energy\\_Sources/Hydro/Hydro\\_Licensing/Klamath\\_River/2014DWF\\_Research\\_Imp\\_Plan\(10-27-14\)F.pdf](http://www.pacificorp.com/content/dam/pacificorp/doc/Energy_Sources/Hydro/Hydro_Licensing/Klamath_River/2014DWF_Research_Imp_Plan(10-27-14)F.pdf)
- David Evans and Associates. 2005. Upper Williamson River Watershed Assessment. Prepared for Klamath Ecosystem Foundation, Upper Williamson River Catchment Group, in Cooperation with the Upper Klamath Basin Working Group and the Klamath Watershed Council by David Evans and Associates, Inc., Portland, OR.  
[http://www.klamathpartnership.org/pdf/uw\\_watershed\\_assessment/uw\\_assessment\\_nomaps.pdf](http://www.klamathpartnership.org/pdf/uw_watershed_assessment/uw_assessment_nomaps.pdf)
- Duff, J.H., K.D. Carpenter, et.al. 2009. Phosphorus and nitrogen legacy in a restoration wetland, upper Klamath Lake, Oregon. *Wetlands*, Vol 29 No 2 pp. 735-746.
- Eldridge, S.L.C., S.A. Wherry, and T.M. Wood. 2014. Statistical analysis of the water-quality monitoring program, Upper Klamath Lake, Oregon, and optimization of the program for 2013 and beyond: U.S. Geological Survey Open File Report 2014-1009, 82 p.,  
<http://dx.doi.org/10.3133/ofr20141009>
- Eilers, J.M., J. Kann, J. Cornett, K. Moser, and A. St. Amand. 2004. Paleolimnological evidence of a change in a shallow, hypereutrophic lake: Upper Klamath Lake, Oregon. *Hydrobiologia* 520: 7-18.
- Federal Energy Regulatory Commission. 2007. Final Environmental Impact statement for the Klamath Hydroelectric Project, Docket No. P-2082-027. 11/18/07. U.S. DOE, FERC, Washington D.C.
- Ferguson, D. and J. Watkins. 2015. Upper Klamath Lake Watershed Ranch Production and Sustainability. Presentation at the Klamath Basin Monitoring Program spring meeting in Yreka, CA, April 29, 2015. Natural Resources Conservation Service and Klamath Soil and Water Conservation District.  
[http://www.kbmp.net/images/stories/pdf/Meeting\\_Materials/Meeting\\_16/6\\_KBMP\\_4-2015.pdf](http://www.kbmp.net/images/stories/pdf/Meeting_Materials/Meeting_16/6_KBMP_4-2015.pdf)
- Fetcho, K. 2006. Klamath River Blue-Green Algae Bloom Report, Water Year 2005. Yurok Tribe Environmental Program, Klamath, CA.
- Friedrichsen, P. T. 1996. Summertime stream temperatures in the North and South Forks of the Sprague River, South Central Oregon. Master's thesis. Department of Forest Engineering, Oregon State University, Corvallis, Oregon.  
<http://ir.library.oregonstate.edu/xmlui/handle/1957/9502>
- Gannett, M.W., K.E. Lite Jr., J.L. La Marche, B.J. Fisher, and D.J. Polette. 2007. Ground-water hydrology of the upper Klamath Basin, Oregon and California: U.S. Geological Survey Scientific Investigations Report 2007-5050, 84 p.
- Genzoli, L. 2013. Shifts in Klamath River metabolism following cyanobacterial bloom. MS thesis. University of Wyoming, Laramie, Wyoming. 53 p.  
<http://search.proquest.com/docview/1494791470>

- Genzoli, L. and R.O. Hall, 2016. Shifts in Klamath River Metabolism Following a Reservoir Cyanobacterial Bloom. *Freshwater Science*:000–000. doi: 10.1086/687752.
- Genzoli, L., R.O. Hall, J.E. Asarian, and J. Kann 2015. Variation and Environmental Association of Ecosystem Metabolism in the Lower Klamath River: 2007-2014. Prepared by the University of Wyoming, Riverbend Sciences, and Aquatic Ecosystem Sciences LLC. for the Klamath Tribal Water Quality Consortium. 44 p. + Appendices.
- Hamilton, J.B., G.L. Curtis, S.M. Snedaker, and D.K. White. 2005. Distribution of Anadromous Fishes in the Upper Klamath River Watershed prior to Hydropower dams—A Synthesis of the Historical Evidence. *Fisheries* 30:10–20.
- Hendrixson, H. 2014. Fourmile Wetland Restoration: One slice at a time. Presentation to Klamath Basin Monitoring Program fall meeting, November 6, 2014 in Yreka, CA. [http://www.kbmp.net/images/stories/pdf/Meeting\\_Materials/Meeting\\_15/23\\_KBMP\\_Fall\\_2014\\_meeting\\_TNC.pdf](http://www.kbmp.net/images/stories/pdf/Meeting_Materials/Meeting_15/23_KBMP_Fall_2014_meeting_TNC.pdf)
- Hendrixson, H. and Bottcher, J. 2015. Recent and Future Restoration in the Upper Klamath Basin. Presentation to Klamath Basin Monitoring Program spring meeting, April 25, 2015 in Yreka, CA. [http://www.kbmp.net/images/stories/pdf/Meeting\\_Materials/Meeting\\_16/22\\_KBMP\\_2015\\_Thursday\\_HENDRIXSONBOTTCHEP.pdf](http://www.kbmp.net/images/stories/pdf/Meeting_Materials/Meeting_16/22_KBMP_2015_Thursday_HENDRIXSONBOTTCHEP.pdf)
- Hoopa Valley Tribal Environmental Protection Agency (HVTEPA). No date. Hoopa Valley Indian Reservation Non-Point Source Pollution Assessment and Management Program. Hoopa Valley Tribal Environmental Protection Agency, Hoopa, CA. 285 p.
- Hoopa Valley Tribal Environmental Protection Agency (HVTEPA). 2008. Water Quality Control Plan Hoopa Valley Indian Reservation. Hoopa Valley Tribal Environmental Protection Agency, Hoopa, CA. 285 p.
- Hoopa Valley Tribal Environmental Protection Agency (HVTEPA). 2013. Water Quality Monitoring by the Hoopa Tribal Environmental Protection Agency 2008-2012 . Prepared by the Hoopa Tribal Environmental Protection Agency in cooperation with Kier Associates. 21 p.
- Horne, A., K. O’Hara, and K. Carlson. 2009. Feasibility and Conceptual Design of an Air Injection Diffuser System in J.C. Boyle Reservoir. Draft Report. Prepared by CH2M HILL and Alex Horne Associates. Prepared for PacifiCorp Energy. October 2009. [http://www.pacificorp.com/content/dam/pacificorp/doc/Energy\\_Sources/Hydro/Hydro\\_Licensing/Klamath\\_River/2008PCKlamathJCBAerationFReport\\_Oct2009.pdf](http://www.pacificorp.com/content/dam/pacificorp/doc/Energy_Sources/Hydro/Hydro_Licensing/Klamath_River/2008PCKlamathJCBAerationFReport_Oct2009.pdf)
- Independent Multidisciplinary Science Team (IMST). 1999. Recovery of Wild Salmonids in Western Oregon Forests: Oregon Forest Practices Act Rules and the Measures in the Oregon Plan for Salmon and Watersheds. Tech. Rpt. 1999-1, Governors Natural Resources Office, Salem OR.
- Jassby, A. and J. Kann. 2010. Upper Klamath Lake monitoring program: preliminary analysis of status and trends for 1990–2009. Prepared for Klamath Tribes Natural Resources Department, Chiloquin, Oregon.
- Kann, J. 1993. Agency Lake Limnology, 1990-91. In: C. Campbell (ed.). Environmental Research in the Klamath Basin, Oregon - 1991 Annual Report. Bureau of Reclamation Technical Report R-93-13. pp. 103-110.

- Kann, J. 1998. Ecology and water quality dynamics of a shallow hypereutrophic lake dominated by Cyanobacteria (*Aphanizomenon flos-aquae*). Doctoral Dissertation. University of North Carolina. Curriculum in Ecology. Chapel Hill, North Carolina.  
[http://cf.unc.edu/ecology/theses/Kahn\\_PhD.pdf](http://cf.unc.edu/ecology/theses/Kahn_PhD.pdf)
- Kann, J. and V.H. Smith. 1999. Estimating the probability of exceeding elevated pH values critical to fish populations in a hypereutrophic lake. *Canadian Journal of Fisheries and Aquatic Sciences* 56(12): 2262-2270.
- Kann, J. and E.B. Welch. 2005. Wind control on water quality in shallow, hypereutrophic Upper Klamath Lake, Oregon. *Lake and Reserv. Manage.* 21: 149–158.
- Kann, J. 2006. *Microcystis aeruginosa* Occurrence in the Klamath River System of Southern Oregon and Northern California. Report for the Yurok Tribe Environmental Program and Fisheries Department, Klamath, CA by Aquatic Ecosystem Sciences, Ashland, OR. 26 p. Accessed online 11/5/2008 at:  
<http://www.yuroktribe.org/departments/ytepd/documents/KannFinalYurokMsaeTechMemo2-3-06.pdf>
- Kann, J. 2008. Technical Memorandum: Microcystin Bioaccumulation in Klamath River Fish and Mussel Tissue: Preliminary 2007 Results. Prepared by Aquatic Ecosystem Sciences, Ashland, OR for the Karuk Tribe Department of Natural Resources, Orleans, CA. 13 pp. + appendices.
- Kann, J. 2010. Compilation of Klamath Tribes Upper Klamath Lake water quality data, 1990–2009. Prepared by Aquatic Ecosystem Sciences, LLC, Ashland, Oregon for the Klamath Tribes Natural Resources Department, Chiloquin, Oregon.
- Kann, J., J.E. Asarian, and A. St. Amand. 2015. Initial Analysis of 1990-2013 Phytoplankton and Zooplankton Data for Upper Klamath Lake (Phase I). Prepared by Aquatic Ecosystem Sciences LLC. for the Klamath Tribes Natural Resources Department. 100p. + appendices.
- Kann, J. and S. Corum. 2006. Summary of 2005 Toxic *Microcystis aeruginosa* Trends in Copco and Iron Gate Reservoirs on the Klamath River, CA. Prepared For: Karuk Tribe Department of Natural Resources, Orleans, CA.
- Kann, J. and S. Corum. 2007. Summary of 2006 Toxic *Microcystis aeruginosa* Trends in Copco and Iron Gate Reservoirs on the Klamath River, CA. Prepared For: Karuk Tribe Department of Natural Resources , Orleans, CA. 23 pp.
- Kann J, S. Corum, and K. Fetcho 2010. Microcystin Bioaccumulation in Klamath River Freshwater Mussel Tissue: 2009 Results. Prepared by Aquatic Ecosystem Sciences, LLC., the Karuk Tribe Natural Resources Department, and the Yurok Tribe Environmental Program:23 pp. + appendices.
- Kann, J. and W.W. Walker. 1999. Nutrient and hydrologic loading to Upper Klamath Lake, Oregon, 1991-1998. Prepared for Klamath Tribes Natural Resource Department, U.S. Bureau of Reclamation Cooperative Studies. 48 p + appendices.
- Karuk Tribe. 2003. Non-Point Source Monitoring Plan for Road Removal Projects. Karuk Tribe of California Department of Natural Resources, Orleans, CA
- Karuk Tribe. 2014. Water Quality Control Plan. Karuk Tribe Department of Natural Resources, Orleans, CA. 45 p.



- Klamath Basin Rangeland Trust (KBRT). 2011. Klamath Basin Rangeland Trust Water Transactions Program. Klamath Basin Rangeland Trust, Klamath Falls, OR.  
[https://www.oregon.gov/owrd/law/docs/GrantApp/GC0013\\_09\\_WTPFinal\\_June2011\\_OWrd.pdf](https://www.oregon.gov/owrd/law/docs/GrantApp/GC0013_09_WTPFinal_June2011_OWrd.pdf)
- Klamath Tribes. 2015. Klamath Tribes History. <http://klamathtribes.org/history>. Accessed 6/13/2016.
- Klamath Watershed Partnership (KWP) and Oregon Department of Environmental Quality (ODEQ). 2015. Project Name: Urban Issues Working Group Non-Point Source Education Project, Final Report. Klamath Watershed Partnership, Klamath Falls, Oregon.
- Klamath Tribal Water Quality Consortium (KTWQC). 2015. Klamath Tribal Water Quality Consortium Strategic Plan and Bylaws.
- Kuwabara, J. S., D. D. Lynch, B. R. Topping, F. Murphy, J. L. Carter, N.S. Simon, F. Parchaso, T. M. Wood, M. K. Lindenberg, K. Wiese, and R. J. Avanzino. 2007. Quantifying the benthic source of nutrients to the water column of Upper Klamath Lake, Oregon. Open-File Report 2007–1276. U. S. Geological Survey, Reston, Virginia.
- Kuwabara, J. S., B.R. Topping, D. D. Lynch, J. L. Carter, and H. I. Essais. 2009. Benthic nutrient sources to hypereutrophic Upper Klamath Lake, Oregon, USA. *Environmental Toxicology and Chemistry* 28: 516–524.
- Kuwabara, J. S., B. R. Topping, J. L. Carter, T. M. Wood, F. Parchaso, J. M. Cameron, J. R. Asbill, R. A. Carlson, and S. V. Fend. 2012. Time scales of change in chemical and biological parameters after engineered levee breaches adjacent to Upper Klamath and Agency Lakes, Oregon. Open-File Report 2012-1057. U. S. Geological Survey.  
<http://pubs.usgs.gov/of/2012/1057>
- Laenen, A., and A. P. LeTourneau. 1996. Upper Klamath Basin nutrient-loading study—estimate of wind-induced resuspension of bed sediment during periods of low lake elevation. Open-File Report 95-414. U. S. Geological Survey, Portland, Oregon.
- Lanman, C.W., K. Lundquist, H. Perryman, J.E. Asarian, B. Dolman, R.B. Lanman, and M.M. Pollock, 2013. The Historical Range of Beaver (*Castor Canadensis*) in Coastal California: An Updated Review of the Evidence. *California Fish and Game* 99:193–221.
- Lindenberg, M.K., Hoilman, G., and T.M. Wood. 2009. Water quality conditions in Upper Klamath and Agency Lakes, Oregon, 2006: U.S. Geological Survey Scientific Investigations Report 2008-5201, 54 p. <http://pubs.usgs.gov/sir/2008/5201/>
- Lyon, S., A. Horne, J. Jordahl, H. Emond, and K. Carlson. 2009. Preliminary Feasibility Assessment of Constructed Treatment Wetlands in the Vicinity of the Klamath Hydroelectric Project. Prepared by CH2M HILL and Alex Horne Associates for PacifiCorp Energy, Portland, Oregon.
- Majerova, M., B.T. Neilson, N.M. Schmadel, J.M. Wheaton, and C.J. Snow. 2015. Impacts of Beaver Dams on Hydrologic and Temperature Regimes in a Mountain Stream. *Hydrol. Earth Syst. Sci.* 19:3541–3556. doi: 10.5194/hess-19-3541-2015.

Mobley Engineering Inc. (MEI). 2008. Reservoir Oxygenation Feasibility Evaluation Report for Copco and Iron Gate Reservoirs. Prepared for PacifiCorp, Portland, OR by Mobley Engineering Inc. with assistance from WolffWare and Reservoir Environmental Management Inc. April 2008. [http://www.pacificorp.com/content/dam/pacificorp/doc/Energy\\_Sources/Hydro/Hydro\\_Licensing/Klamath\\_River/Reservoir\\_Oxygenation\\_Feasibility\\_Evaluation\\_Report\\_Iron\\_Gate\\_Copco\\_Reservoirs.pdf](http://www.pacificorp.com/content/dam/pacificorp/doc/Energy_Sources/Hydro/Hydro_Licensing/Klamath_River/Reservoir_Oxygenation_Feasibility_Evaluation_Report_Iron_Gate_Copco_Reservoirs.pdf)

Morace, J.L. 2007. Relation between selected water-quality variables, climatic factors, and lake levels in Upper Klamath and Agency Lakes, Oregon, 1990–2006. U.S. Geological Survey Scientific Investigations Report 2007-5117. <http://pubs.water.usgs.gov/sir20075117>.

National Marine Fisheries Service (NMFS). 2014. Final Recovery Plan for the Southern Oregon/Northern California Coast Evolutionarily Significant Unit of Coho Salmon (*Oncorhynchus kisutch*). National Marine Fisheries Service. Arcata, CA. [http://www.westcoast.fisheries.noaa.gov/protected\\_species/salmon\\_steelhead/recovery\\_planning\\_and\\_implementation/southern\\_oregon\\_northern\\_california\\_coast/southern\\_oregon\\_northern\\_california\\_coast\\_salmon\\_recovery\\_domain.html](http://www.westcoast.fisheries.noaa.gov/protected_species/salmon_steelhead/recovery_planning_and_implementation/southern_oregon_northern_california_coast/southern_oregon_northern_california_coast_salmon_recovery_domain.html)

National Research Council (NRC). 2004. Endangered and Threatened Fishes in the Klamath River Basin: Causes of Decline and Strategies for Recovery. Washington, DC: The National Academies Press.

Natural Resources Conservation Service. 2009. Sprague River CEAP Study Report. U.S. Department of Agriculture, Oregon Natural Resources Conservation Service. [http://www.nrcs.usda.gov/Internet/FSE\\_DOCUMENTS/nrcseprd357894.pdf](http://www.nrcs.usda.gov/Internet/FSE_DOCUMENTS/nrcseprd357894.pdf).

Natural Resources Conservation Service (NRCS). 2010. Wood River, Upper Klamath Basin, Oregon Conservation Effects Assessment Project, Special Emphasis Watershed, Final Report. U.S. Department of Agriculture, Oregon Natural Resources Conservation Service. [http://www.nrcs.usda.gov/Internet/FSE\\_DOCUMENTS/nrcseprd334614.pdf](http://www.nrcs.usda.gov/Internet/FSE_DOCUMENTS/nrcseprd334614.pdf).

NewFields River Basin Services and G. M. Kondolf. 2012. Evaluating Stream Restoration Projects in the Sprague River Basin. Prepared for the Klamath Watershed Partnership in conjunction with the Klamath Tribes, the U.S. Fish and Wildlife Service, the Klamath Basin Rangeland Trust, Sustainable Northwest, and The Nature Conservancy.

North Coast Regional Water Quality Control Board (NCRWQCB). 2010. Final Staff Report for the Klamath River Total Maximum Daily Loads (TMDLs) addressing temperature, dissolved oxygen, nutrient, and microcystin impairments in California, the proposed site specific dissolved oxygen objectives for the Klamath River in California, and the Klamath River and Lost River implementation plans. State of California.

North Coast Regional Water Quality Control Board (NCRWQCB). 2011. Water Quality Control Plan for the North Coast Region. North Coast Regional Water Quality Control Board, Santa Rosa, CA. 274 p.

North Coast Regional Water Quality Control Board (NCRWQCB). 2015. Order No. R1-2015-0021, Waiver of Waste Discharge Requirements for Nonpoint Source Discharges Related to Certain Federal Land Management Activities on National Forest System Lands in the North Coast Region. California Regional Water Quality Control Board, North Coast Region, Santa Rosa, CA. [http://www.waterboards.ca.gov/northcoast/board\\_decisions/adopted\\_orders/pdf/2015/15\\_0021\\_Waiver\\_USFS.pdf](http://www.waterboards.ca.gov/northcoast/board_decisions/adopted_orders/pdf/2015/15_0021_Waiver_USFS.pdf)

O'Connor, J.E., P.F. McDowell, P. Lind, C.G. Rasmussen, and M.K. Keith. 2015. Geomorphology and flood-plain vegetation of the Sprague and lower Sycan Rivers, Klamath Basin, Oregon: U.S. Geological Survey Scientific Investigations Report 2014-5223, 122 p., 1 pl., <http://dx.doi.org/10.3133/sir20145223>.

Office of Environmental Health Hazard Assessment (OEHHA). 2005. Memo from Dr. Karlyn Kaley, EPA Toxicologist, to Matt St. John, North Coast Regional Water Quality Control Board, re: Cyanobacterial Microcystin Toxin Summer 2005 Water Sampling Results from Copco and Iron Gate Reservoirs. Integrated Risk Assessment Branch, Office of Environmental Health Hazard Assessment, California Environmental Protection Agency, Sacramento, CA 4 p.

Office of Environmental Health Hazard Assessment (OEHHA). 2008 . Information Related to the Occurrence of Microcystin in the Tissues of Klamath River Biota .Letter from George Alexeeff to OEHHA to Randy Landolt of PacifiCorp, August 6,2008. OEHHA, Sacramento, CA. 5pp.

Office of Environmental Health Hazard Assessment (OEHHA). 2012. Toxicological Summary and Suggested Action Levels to Reduce Potential Adverse Health Effects of Six Cyanotoxins. Final Report-- May 2012. Office of Environmental Health Hazard Assessment California Environmental Protection Agency, Sacramento, California 95812-4010. [http://www.waterboards.ca.gov/water\\_issues/programs/peer\\_review/docs/calif\\_cyanotoxins/cyanotoxins053112.pdf](http://www.waterboards.ca.gov/water_issues/programs/peer_review/docs/calif_cyanotoxins/cyanotoxins053112.pdf)

Oregon Department of Agriculture (ODA). 2011. Klamath Headwaters Agricultural Water Quality Management Area Plan. Prepared with assistance from the Klamath Headwaters Local Advisory Committee and the Klamath Soil and Water Conservation District. ODA, Portland, OR. 55 p. <https://www.oregon.gov/ODA/shared/Documents/Publications/NaturalResources/KlamathAWQMAreaPlan.pdf>

Oregon Department of Agriculture (ODA) and Lost River Local Agricultural Water Quality Advisory Committee (LRLAWQAC). 2015. Lost River Agricultural Water Quality Management Area Plan. Prepared with assistance from the Klamath Headwaters Local Advisory Committee and the Klamath Soil and Water Conservation District. ODA, Portland, OR. 60 p. <https://www.oregon.gov/ODA/shared/Documents/Publications/NaturalResources/LostRiverAWQMAreaPlan.pdf>

Oregon Department of Environmental Quality (ODEQ). 2002. Upper Klamath Lake Drainage Total Maximum Daily Load (TMDL) and Water Quality Management Plan (WQMP). Oregon Department of Environmental Quality. Portland, Oregon. 204 p.

Oregon Department of Environmental Quality (ODEQ). 2005. Table 180A Designated Beneficial Uses Klamath Basin (340-41-0180). Portland, Oregon. <http://www.deq.state.or.us/wq/rules/div041/dbutables/table180a.pdf>.

Oregon Department of Environmental Quality (ODEQ). 2010a. Upper Klamath and Lost River Subbasins TMDL, Appendix A: Temperature Model Calibration Report. ODEQ 10-WQ-030. Portland, Oregon. <http://www.deq.state.or.us/wq/tmdls/klamath.htm#upks>

Oregon Department of Environmental Quality (ODEQ). 2010b. Upper Klamath and Lost River Subbasins Total Maximum Daily Load (TMDL) and Water Quality Management Plan (WPMP). ODEQ 10-WQ-030. Portland, Oregon. <http://www.deq.state.or.us/wq/tmdls/klamath.htm#upks>

Oregon Department of Environmental Quality (ODEQ). 2012. Methodology for Oregon's 2012 Water Quality Report and List of Water Quality Limited Waters. State of Oregon Department of Environmental Quality, Portland, OR. 127 p.

Oregon Watershed Enhancement Board (OWEB). 2016. Upper Klamath Special Investment Partnership, Accomplishments Summary Report, 2012-2015. Oregon Watershed Enhancement Board, Salem, OR. 10 p. <http://www.oregon.gov/OWEB/docs/board/2016/April/UKSIP-Accomplishments-Report-2016.pdf>

Otten, T.G, J.R. Crosswell, S. Mackey, and T W. Dreher. 2015. Application of molecular tools for microbial source tracking and public health risk assessment of a *Microcystis* bloom traversing 300 km of the Klamath River. *Harmful Algae* 46:71-81

PacifiCorp. 2004. Final License Application for the Klamath River Hydroelectric Project (FERC Project No. 2082). Portland, OR.

PacifiCorp, 2005. Response to FERC AIR AR-2, Final Technical Report, Anadromous Fish Restoration, Klamath Hydroelectric Project (FERC Project No. 2082). PacifiCorp, Portland, OR

PacifiCorp. 2014. Application for Water Quality Certification Pursuant to Section 401 of the Federal Clean Water Act for the Relicensing of the Klamath Hydroelectric Project (FERC No. 2082) in Siskiyou County, California, Klamath Hydroelectric Project. (FERC Project No. 2082). PacifiCorp, Portland, OR

PacifiCorp. 2015. Study Plan Klamath Hydroelectric Project Interim Measure 11 Study Activities for 2015, May 29, 2015.

Pischel, E.M., and M.W. Gannett. 2015. Effects of groundwater pumping on agricultural drains in the Tule Lake subbasin, Oregon and California: U.S. Geological Survey Scientific Investigations Report 2015-5087, 44 p. <http://dx.doi.org/10.3133/sir20155087>.

Pollock, M.M., G.R. Pess, T.J. Beechie, and D.R. Montgomery. 2004. The Importance of Beaver Ponds to Coho Salmon Production in the Stillaguamish River Basin, Washington, USA. *North American Journal of Fisheries Management* 24:749-760.

Pollock, M.M., T.J. Beechie, J.M. Wheaton, C.E. Jordan, N. Bouwes, N. Weber, and C. Volk. 2014. Using Beaver Dams to Restore Incised Stream Ecosystems. *BioScience* 64:279-290. doi: 10.1093/biosci/biu036.

Pollock, M.M., G. Lewallen, K. Woodruff, C.E. Jordan and J.M. Castro (Editors). 2015. The Beaver Restoration Guidebook: Working with Beaver to Restore Streams, Wetlands, and Floodplains. Version 1.02. United States Fish and Wildlife Service, Portland, Oregon. 189 pp. <http://www.fws.gov/oregonfwo/ToolsForLandowners/RiverScience/Beaver.asp>

Quartz Valley Indian Reservation (QVIR). 2008. Quartz Valley Indian Reservation Watershed Based Non-Point Source Management Plan, Quartz Valley, California. QVIR Environmental Department, Fort Jones, CA. 45 p.

Quartz Valley Indian Reservation. 2016. Quality Assurance Project Plan for Water Quality Sampling and Analysis, CWA 106 grant identification # I-96927206-0. Prepared by Quartz Valley Indian Reservation, Fort Jones, CA.

Resighini Rancheria Environmental Protection Agency (REPA). 2006. The Resighini Rancheria Tribal Water Quality Ordinance Number 01-2002. Resighini Rancheria Environmental Protection Agency, Klamath CA. 43 p.

Resighini Rancheria Environmental Protection Agency (REPA). 2010. Surface Water Sampling and Analysis Plan, Resighini Rancheria, Section 106 Water Quality Assessment, Clean Water Act Project. Resighini Rancheria Environmental Protection Agency, Klamath CA. 18 p.

Risley, J. C., and A. Laenen. 1999. Upper Klamath Lake Basin nutrient-loading study—assessment of historic flows in the Williamson and Sprague rivers. Water-Resources Investigations Report 98-4198. U. S. Geological Survey, in cooperation with U. S. Bureau of Reclamation, Portland, Oregon.

Schwankl, L.J., Prichard, T.L., and B.R. Hanson. 2007. Tailwater Return Systems. University of California Division of Agriculture and Natural Resources, Publication 8225, 10 p.  
<http://anrcatalog.ucanr.edu/pdf/8225.pdf>

Scott, N. 2016. Treatment Wetland Pilot Project Development in the Upper Klamath Basin. Presentation to Klamath Basin Monitoring Program spring meeting, April 21, 2016 in Klamath Falls, OR.  
[http://www.kbmp.net/images/stories/pdf/Meeting\\_Materials/Meeting\\_18/14\\_KBMP\\_Spring\\_2016.pdf](http://www.kbmp.net/images/stories/pdf/Meeting_Materials/Meeting_18/14_KBMP_Spring_2016.pdf)

Shasta Valley Resource Conservation District (SVRCD). No date. Irrigation Tailwater Management in the Shasta Valley (brochure). Shasta Valley Resource Conservation District, Yreka, CA. <http://svrcd.org/wordpress/wp-content/uploads/2012/01/Final-tailwatrpub.pdf>

Silloway, S. 2010. Fish Surveys Related to the Proposed Del Norte Highway 101 Klamath Grade Raise Project Contract No. 03A1317 Task Order 48.  
<http://www.yuroktribe.org/departments/fisheries/documents/YTFP2010KlamathGradeRaisebyCaltransFishReportFINAL.pdf>

Simon, N. S., and S. N. Ingle. 2011. Physical and chemical characteristics including total and geochemical forms of phosphorus in sediment from the top 30 centimeters of cores collected in October 2006 at 26 sites in Upper Klamath Lake, Oregon. Open-File Report 2011-1168. U. S. Geological Survey. <http://pubs.usgs.gov/of/2011/1168>

Simon, N. S., D. Lynch, and T. N. Gallaher. 2009. Phosphorus fractionation in sediment cores collected in 2005 before and after onset of an *Aphanizomenon flos-aquae* bloom in Upper Klamath Lake, OR, USA. *Water Air Soil Pollution*. 204: 139–153.

Smith, J.E., and K.A. Rykbost. 2000. Klamath Basin Crop Trends. Klamath Experiment Station, Klamath Falls, OR.  
<http://oregonstate.edu/dept/kbrec/sites/default/files/documents/ag/ar00chpt1.pdf>

Snyder, D.T. and J.L. Morace. 1997. Nitrogen and phosphorus loading from drained wetlands adjacent to Upper Klamath and Agency Lakes, Oregon: U.S. Geological Survey Water Resources Investigations Report 97-4059. 67 p.

State Water Resources Control Board (SWRCB). 2015a. California Nonpoint Source Program Implementation Plan 2014 - 2020. State Water Resources Control Board, Sacramento, California.  
[http://www.waterboards.ca.gov/water\\_issues/programs/nps/docs/plans\\_policies/sip\\_2014to2020.pdf](http://www.waterboards.ca.gov/water_issues/programs/nps/docs/plans_policies/sip_2014to2020.pdf)

State Water Resources Control Board (SWRCB). 2015b. Staff Report 2012 California Integrated Report Clean Water Act Sections 303(d) and 305(b). California Environmental Protection

- Agency, State Water Resources Control Board, Sacramento, California.  
[http://www.waterboards.ca.gov/water\\_issues/programs/tmdl/integrated2012.shtml](http://www.waterboards.ca.gov/water_issues/programs/tmdl/integrated2012.shtml)
- Strecker, E.W. 2001. Low-Impact Development (LID)—Is It Really Low or Just Lower? Linking Stormwater BMP Designs and Performance to Receiving Water Impact Mitigation. ASCE, pp. 210–222.
- Stillwater Sciences, Riverbend Sciences, Aquatic Ecosystem Sciences, Atkins, Tetra Tech, NSI/Biohabitats, and Jones & Trimiew Design. 2012. Klamath River pollutant reduction workshop—information packet. Revised. Prepared for California State Coastal Conservancy, Oakland, California.
- Stillwater Sciences, Jones & Trimiew Design, Atkins, Tetra Tech, Riverbend Sciences, Aquatic Ecosystem Sciences, and NSI/Biohabitats. 2013. Water Quality Improvement Techniques for the Upper Klamath Basin: A Technical Workshop and Project Conceptual Designs. Prepared for California State Coastal Conservancy, Oakland, California. 106 p.
- Sullivan, A.B., S.A. Rounds, M.L. Deas, J.R. Asbill, R.E. Wellman, M.A. Stewart, M.W. Johnston, and I.E. Sogutlugil. 2011. Modeling Hydrodynamics, Water Temperature, and Water Quality in the Klamath River Upstream of Keno Dam, Oregon, 2006–09. U.S. Geological Survey Scientific Investigations Report 2011-5105. <http://pubs.usgs.gov/sir/2011/5105/>.
- Sullivan, A.B., I.E. Sogutlugil, M.L. Deas, and S.A. Rounds. 2014. Water-quality modeling of Klamath Straits Drain recirculation, a Klamath River wetland, and 2011 conditions for the Link River to Keno Dam reach of the Klamath River, Oregon: U.S. Geological Survey Open-File Report 2014–1185, 75 p., <http://dx.doi.org/10.3133/ofr20141185>.
- Sullivan, A.B., I.E. Sogutlugil, S.A. Rounds, and M.L. Deas. 2013. Modeling the water-quality effects of changes to the Klamath River upstream of Keno Dam, Oregon: U.S. Geological Survey Scientific Investigations Report 2013–5135, 60 p., <http://pubs.usgs.gov/sir/2013/5135>.
- Timmons, D. 2016. Water Rights Settlements in Oregon’s Klamath Basin Facing Uncertain Future. <http://www.martenlaw.com/newsletter/20160222-water-rights-klamath-basin>. Accessed 6/13/2016
- Upper Klamath Basin Comprehensive Agreement (UKBCA). 2014. Proposed Upper Klamath Basin comprehensive agreement. <http://klamathtribes.org/wp-content/uploads/2014/08/2014-4-18-UPPER-KLAMATH-BASIN-COMPREHENSIVE-AGREEMENT.pdf>. Accessed 6/13/2016.
- U.S. Bureau of Reclamation (USBR). 2005. Natural flow of the upper Klamath River-Phase I. Natural inflow to, natural losses from, and natural outfall of Upper Klamath Lake to the Link River and the Klamath River at Keno. USBR Technical Service Center, Klamath Basin Area Office, Klamath Falls, OR
- U.S. Bureau of Reclamation (USBR). 2016a. Klamath Project. [http://www.usbr.gov/projects/Project.jsp?proj\\_Name=Klamath+Project](http://www.usbr.gov/projects/Project.jsp?proj_Name=Klamath+Project). Accessed Jan 28, 2016.
- U.S. Bureau of Reclamation (USBR). 2016b. Klamath Project Water Quality and Use – Initial Demonstration Assessment, Upper Klamath Basin, Oregon. Prepared by the Bureau of Reclamation Technical Service Center in Denver, Colorado and Klamath Basin Area Office in Klamath Falls, Oregon. 126 p., plus appendices.

U.S. Department of Agriculture (USDA). 2009. Sprague River conservation effects assessment project (CEAP) study report: U.S. Department of Agriculture, Natural Resources Conservation Service, 100 p. [http://www.nrcs.usda.gov/Internet/FSE\\_DOCUMENTS/nrcseprd357894.pdf](http://www.nrcs.usda.gov/Internet/FSE_DOCUMENTS/nrcseprd357894.pdf)

U.S. Department of Agriculture (USDA). 2016a. Klamath: Taking on drought before it occurs. United State Department of Agriculture, Natural Resources Conservation District - Oregon. <http://www.nrcs.usda.gov/wps/portal/nrcs/detail/or/programs/?cid=nrcseprd429622>

U.S. Department of Agriculture (USDA). 2016b. Regional Conservation Partnership Program (RCPP), Investing in Oregon – 2016. United State Department of Agriculture. [http://www.nrcs.usda.gov/wps/PA\\_NRCSCconsumption/download?cid=nrcseprd630007&ext=pdf](http://www.nrcs.usda.gov/wps/PA_NRCSCconsumption/download?cid=nrcseprd630007&ext=pdf)

U.S. Department of Interior (US DOI) and California Department of Fish and Game (CDFG). 2012. Klamath Facilities Removal, Environmental Impact Statement/ Environmental Impact Report (EIS/EIR). State Clearinghouse # 2010062060. Sacramento, California. <http://klamathrestoration.gov/Draft-EIS-EIR/download-draft-eis-eir>.

U.S. Environmental Protection Agency (USEPA). 2005. Guidance for 2002 Assessment, listing and reporting requirements pursuant to Sections 303(d) and 305(b) of the Clean Water Act. United States Environmental Protections Agency, Washington D.C.

U.S. Environmental Protection Agency (USEPA). <http://www.epa.gov/wqs-tech/water-quality-standards-regulations-hoopa-valley-tribe> . Accessed Mar 2 2016.

U.S. Environmental Protection Agency. 2008. Lost River, California total maximum daily loads; nitrogen and biochemical oxygen demand to address dissolved oxygen and pH impairments. Final Report. U.S. Environmental Protection Agency, Region IX. <https://www3.epa.gov/region9/water/tmdl/lost-river/TmdlLostRiver12-30-08.pdf>, accessed June 20, 2016.

U.S. Environmental Protection Agency. 2010. Handbook for Developing and Managing Tribal Nonpoint Source Pollution Programs Under Section 319 of the Clean Water Act, EPA 841-B-10-001. U.S. Environmental Protection Agency, Office of Water. <https://www.epa.gov/polluted-runoff-nonpoint-source-pollution/nonpoint-source-tribal-resources-and-training>

U.S. Fish and Wildlife Service (USFWS). 2001. Biological/Conference Opinion Regarding the Effects of Operation of the Bureau of Reclamation's Klamath Project on the Endangered Lost River Sucker (*Deltistes luxatus*), Endangered Shortnose Sucker (*Chasmistes brevirostris*), Threatened Bald Eagle (*Haliaeetus leucocephalus*), and Proposed Critical Habitat for the Lost River/Shortnose Suckers. U.S. Fish and Wildlife Service, Klamath Falls Fish and Wildlife Office, Klamath Falls, OR.

U.S. Fish and Wildlife Service (USFWS). 2003a Klamath River Fish Die-off, September 2002, Report on Estimates of Mortality. US Fish and Wildlife Services Report Number AFWO-01-03.

U.S. Fish and Wildlife Service (USFWS). 2003b Klamath River Fish Die-off, September 2002, Causative Factors of Mortality. US Fish and Wildlife Services Report Number AFWO-02-03.

U.S. Fish and Wildlife Service (USFWS). 2010. Klamath Marsh National Wildlife Refuge Comprehensive Conservation Plan. U.S. Fish and Wildlife Service, Sacramento, California and Tule Lake, California. <http://www.fws.gov/klamathbasinrefuges/KlamathMarshCCP/kmarshccpFINAL.html>

U.S. Fish and Wildlife Service (USFWS). 2012. Revised recovery plan for the Lost River sucker (*Deltistes luxatus*) and shortnose sucker (*Chasmistes brevirostris*). U.S. Fish and Wildlife Service, Pacific Southwest Region, Sacramento, California. xviii + 122 pp.  
[http://www.fws.gov/klamathfallsfwo/suckers/sucker\\_news/FinalRevLRS-SNSRecvPln/FINAL%20Revised%20LRS%20SNS%20Recovery%20Plan.pdf](http://www.fws.gov/klamathfallsfwo/suckers/sucker_news/FinalRevLRS-SNSRecvPln/FINAL%20Revised%20LRS%20SNS%20Recovery%20Plan.pdf).

U.S. Fish and Wildlife Service (USFWS). 2013a. Tools for Landowners, Partners for Fish & Wildlife Program. U.S. Fish and Wildlife Service, Oregon Fish and Wildlife Office.  
<https://www.fws.gov/oregonfwo/toolsforlandowners/partners/>

U.S. Fish and Wildlife Service (USFWS). 2013b. Walking Wetlands. U.S. Fish and Wildlife Service, Tule Lake National Wildlife Refuge. Accessed June 21, 2016:  
<https://www.fws.gov/refuge/tulelake/walkingwetlands.html>

U.S. Forest Service (USFS) and Bureau of Land Management (BLM). 1994. Record of Decision for Amendments to Forest Service and Bureau of Land Management Planning Documents Within the Range of the Northern Spotted Owl: Standards and Guidelines for Management of Habitat for Late-Successional and Old-Growth Forest Related Species Within the Range of the Northern Spotted Owl. USDA Forest Service and USDI Bureau of Land Management, Washington, DC.

Walker, W.W., J.D. Walker, and J. Kann. 2012. Evaluation of water and nutrient balances for the Upper Klamath Lake Basin in water years 1992–2010: Technical Report to the Klamath Tribes Natural Resources Department, 50 p. plus appendixes.  
[http://wwwwalker.net/ukl/klamath\\_nutrientbudget\\_2012\\_final.pdf](http://wwwwalker.net/ukl/klamath_nutrientbudget_2012_final.pdf)

Walker, J. D., J. Kann, and W.W. Walker. 2015. Spatial and temporal nutrient loading dynamics in the Sprague River Basin, Oregon. Prepared by Aquatic Ecosystem Sciences, J. D. Walker, and W. W. Walker for the Klamath Tribes Natural Resources Department. 73p. + appendices.  
[https://s3.amazonaws.com/walkerenvres.com/reports/klamath/2015-sprague-nutrient-dynamics/Sprague\\_River\\_Nutrient\\_Dynamics\\_20151229\\_final.pdf](https://s3.amazonaws.com/walkerenvres.com/reports/klamath/2015-sprague-nutrient-dynamics/Sprague_River_Nutrient_Dynamics_20151229_final.pdf)

Watercourse Engineering, Inc. 2006. Characterization of Organic Matter Fate and Transport in the Klamath River below Link Dam to Assess Treatment/Reduction Potential. Prepared for U.S. Bureau of Reclamation. Davis, CA. 167 p.

Watercourse Engineering, Inc. 2013. Evaluation of Particulate Organic Matter Removal from Klamath River Source Water Using Stormwater Treatment Technology, 2012. Prepared for PacifiCorp Energy, Portland OR. April.

Watercourse Engineering, Inc. 2014a. Draft Technical Report, Conceptual Design Evaluation for Full-Scale Particulate Organic Matter Removal from Klamath River Source Water Using Stormwater Treatment Technology, 2013. Prepared for PacifiCorp Energy, Portland OR. June.

Watercourse Engineering, Inc. 2014b. Evaluation of Particulate Organic Matter Removal from Klamath River Source Water Using Stormwater Treatment Technology, 2013. Prepared for PacifiCorp Energy, Portland OR. June.

Watercourse Engineering, Inc. 2015. Klamath River Baseline Water Quality Sampling, 2014 Annual Report. Prepared for the KHSWA Water Quality Monitoring Group by Watercourse Engineering, Davis, CA.  
[http://www.pacificorp.com/content/dam/pacificorp/doc/Energy\\_Sources/Hydro/Hydro\\_Licensin g/Klamath\\_River/2015-IM15-Study-Plan-final.pdf](http://www.pacificorp.com/content/dam/pacificorp/doc/Energy_Sources/Hydro/Hydro_Licensin g/Klamath_River/2015-IM15-Study-Plan-final.pdf).



Weddell, B.J. 2000. Relationship between Flows in the Klamath River and Lower Klamath Lake prior to 1910. Prepared for the US Department of the Interior, US Fish and Wildlife Service, Klamath Basin Refuges, Tulelake, CA.

Wherry, S.A., T.M. Wood, and C.W. Anderson. 2015. Revision and Proposed Modification for a Total Maximum Daily Load Model for Upper Klamath Lake, Oregon. USGS Numbered Series, U.S. Geological Survey, Reston, VA.

Wood, T.M., R.T. Cheng, J.W. Gartner, G.R. Hoilman, M.K. Lindenberg, and R.E. Wellman. 2008. Modeling hydrodynamics and heat transport in Upper Klamath Lake, Oregon, and implications for water quality: U.S. Geological Survey Scientific Investigations Report 2008–5076, 48 p.

Wood, T.M., H.A. Hendrixson, D.F. Markle, C.S. Erdman, S.M. Burdick, and C.M. Ellsworth. 2014. Simulation and validation of larval sucker dispersal and retention through the restored Williamson River Delta and Upper Klamath Lake system, Oregon: U.S. Geological Survey Scientific Investigations Report 2013–5194, 34 p., <http://dx.doi.org/10.3133/sir20135194>.

Wood, T.M., S.A. Wherry, J.L. Carter, J.S. Kuwabara, N.S. Simon, and S.A. Rounds. 2013. Technical evaluation of a total maximum daily load model for Upper Klamath and Agency Lakes, Oregon: U.S. Geological Survey Open-File Report 2013–1262, 75 p, <http://dx.doi.org/10.3133/ofr20131262>

Wong, S., and H. Hendrickson. 2011. Water quality conditions in the Williamson River Delta, Oregon: three years post-restoration. 2010 Annual Data Report. The Nature Conservancy Klamath Basin Field Office, Klamath Falls, OR.

Yurok Tribe Environmental Program (YTEP). 2004. Water Quality Control Plan for the Yurok Indian Reservation. Yurok Tribe Environmental Program, Klamath, CA. 37 p.

Yurok Tribe Environmental Program (YTEP). 2014. NPS Assessment and Management Program Plan. Yurok Tribe Environmental Program, Klamath, CA. 147 p.

## 13 ACRONYMS AND ABBREVIATIONS LIST

AGR	Agriculture
Al	Aluminum
AMPP	Assessment and Management Program Plan
BDAs	Beaver Dam Analogues
BLM	United States Bureau of Land Management
BMPs	Best Management Practices
CCC	California Coastal Conservancy
CDFW	California Department of Fish and Wildlife
CDFG	California Department of Fish and Game
Consortium	Klamath Tribal Water Quality Consortium
CWA	Clean Water Act
DO	Dissolved Oxygen
DSTWs	Diffuse Source Treatment Wetlands
FERC	Federal Energy Regulatory Commission
FFOD	Findings of Fact and Order of Determination
FOR	Forestry
GIS	geographic information system
HU	Hydrologic Unit
HUC	USGS hydrologic unit code
HVTEPA	Hoopa Valley Tribal Environmental Protection Agency
HHA	Hydromodification and Habitat Alteration
IM	Interim Measures
IMST	Independent Multidisciplinary Science Team
KBMP	Klamath Basin Monitoring Program
KBRA	Klamath Basin Restoration Agreement
KHP	Klamath Hydroelectric Project
KHSA	Klamath Hydroelectric Settlement Agreement
KIP	Klamath Irrigation Project
KBREC	Klamath Basin Research & Extension Center
KSWCD	Klamath Soil and Water Conservation District
KTAP	Klamath Tracking and Accounting Program
KTWQC	Klamath Tribal Water Quality Consortium
KWP	Klamath Watershed Partnership
LID	Lower Impact Development
LKL	Lower Klamath Lake
LRLAWQAC	Lost River Local Agricultural Water Quality Advisory Committee
m	meters
MEI	Mobley Engineering Inc.
mi	miles
NCRWQCB	North Coast Regional Water Quality Control Board
NOAA	National Oceanic and Atmospheric Administration

NMFS	National Marine Fisheries Service
NPS	Nonpoint Source
NPS AMPP	Nonpoint Source Assessment and Management Program Plan
NRC	Natural Research Council
NRCS	Natural Resources Conservation Service
NWFP	Northwest Forest Plan
ODA	Oregon Department of Agriculture
ODEQ	Oregon Department of Environmental Quality
OEHHA	Office of Environmental Health Hazard Assessment
OTH	Other
OWEB	Oregon Watershed Enhancement Board
OWRD	Oregon Water Resources Department
PCSRF	Pacific Coastal Salmon Recovery Fund
QVIR	Quartz Valley Indian Reservation
RCPP	Regional Conservation Partnership Program
REPA	Resighini Rancheria Environmental Protection Authority
RIT	Recovery Implementation Team
RM	River Mile
SIF	Specified Instream Flows
SIP	Special Investment Partnership
SVRCD	Shasta Valley Resource Conservation District
SWRCB	State Water Resources Control Board
TAS	treatment as a state
TMDLs	Total Maximum Daily Loads
Ti	Titanium
TU	Trout Unlimited
U.S. BLM	United States Bureau of Land Management
U.S. DOI	United States Department of Interior
UKBCA	Upper Klamath Basin Comprehensive Agreement
UKCAN	Upper Klamath Conservation Action Network
UKL	Upper Klamath Lake
USBR	United States Bureau of Reclamation
USEPA	United States Environmental Protection Agency
USDA	United States Department of Agriculture
USFS	U.S. Forest Service
USFWS	United States Fish and Wildlife Service
USGS	U.S. Geological Survey
WDR	Waste Discharge Requirements
WRP	Wetlands Reserve Program
WUP	Water Use Program
YTEP	Yurok Tribe Environmental Program

## APPENDIX A: OREGON DEPARTMENT OF ENVIRONMENTAL QUALITY LIST OF IMPAIRED WATERBODIES

Table A10. Oregon Department of Environmental Quality (ODEQ) list of waterbodies in the Upper Klamath Basin designated as impaired under section 303(d) of the Clean Water Act. Key to abbreviations: R. = River, Cr. = Creek, S.F. = South Fork, N.F. = North Fork, UKL = Upper Klamath Lake.

Watershed (USGS 4th Field Name)	Water Body (Stream/Lake)	River Miles	Parameter	Season	Criteria	Beneficial Uses	Integrated Category
Lost	Klamath R.	250 to 251	Fecal Coliform	Summer	Fecal coliform log mean of 200 organisms per 100 ml; no more than 10% > 400 per 100 ml	Water contact recreation	Attaining
Lost	Klamath R.	250 to 251	Fecal Coliform	FallWinterSpring	Fecal coliform log mean of 200 organisms per 100 ml; no more than 10% > 400 per 100 ml	Water contact recreation	Attaining
Lost	Klamath R.	250 to 251	Flow Modification	Undefined	Creation of foul tastes, odors, or toxins. <sup>1</sup>	Resident fish and aquatic life; Salmonid fish rearing; 4B Salmonid fish spawning	
Lost	Klamath R.	250 to 251	Nutrients	Undefined		Aesthetics	3
Lost	Klamath R.	250 to 251	pH	FallWinterSpring	pH 6.5 to 9.0	Anadromous fish passage; Resident fish and aquatic life; Salmonid fish rearing; Salmonid fish spawning; Water contact recreation	Attaining
Lost	Klamath R.	250 to 251	Sedimentation	Undefined	Bottom Deposits <sup>2</sup>	Resident fish and aquatic life; Salmonid fish rearing; 3 Salmonid fish spawning	
Lost	Klamath R.	250 to 251	Temperature	Summer	Rearing: 17.8 C	Anadromous fish passage; Salmonid fish rearing	Criteria change or use clarification
Lost	Klamath Strait	0 to 9.8	Alkalinity	Year Around	Table 20 Toxic Substances	Aquatic life	2
Lost	Klamath Strait	0 to 9.8	Ammonia	Year Around	Table 20 Toxic Substances	Aquatic life	5
Lost	Klamath Strait	0 to 9.8	Arsenic	Year Around	Table 40 Human Health Criteria for Toxic Pollutants	Aquatic life; Human health	5
Lost	Klamath Strait	0 to 9.8	Chlorophyll a	FallWinterSpring	Reservoir, river, estuary, non-thermally stratified lake: 0.015 mg/l	Aesthetics; Fishing; Livestock watering; Water contact recreation; Water supply	3
Lost	Klamath Strait	0 to 9.8	Chlorophyll a	Summer	Reservoir, river, estuary, non-thermally stratified lake: 0.015 mg/l	Aesthetics; Fishing; Livestock watering; Water contact recreation; Water supply	5
Lost	Klamath Strait	0 to 9.8	Dissolved Oxygen	Year Around (Non-spawning)	Cool water: Not less than 6.5 mg/l		5
Lost	Klamath Strait	0 to 9.8	Dissolved Oxygen	Year Around	Spawning: Not less than 11.0 mg/L or 95% of saturation	Salmonid fish spawning	Criteria change or use clarification
Lost	Klamath Strait	0 to 9.8	E. Coli	Summer	E. Coli <sup>3</sup>		5
Lost	Klamath Strait	0 to 9.8	E. Coli	Summer	30-day log mean of 126 E. coli	Water contact recreation	2

Watershed (USGS 4th Field Name)	Water Body (Stream/Lake)	River Miles	Parameter	Season	Criteria	Beneficial Uses	Integrated Category
					organisms per 100 ml; no single sample > 406 organisms per 100 ml		
Lost	Klamath Strait	0 to 9.8	E. Coli	FallWinterSpring	30-day log mean of 126 E. coli organisms per 100 ml; no single sample > 406 organisms per 100 ml	Water contact recreation	2
Lost	Klamath Strait	0 to 9.8	Fecal Coliform	Summer	Fecal coliform log mean of 200 organisms per 100 ml; no more than 10% > 400 per 100 ml	Water contact recreation	Criteria change or use clarification
Lost	Klamath Strait	0 to 9.8	Fecal Coliform	FallWinterSpring	Fecal coliform log mean of 200 organisms per 100 ml; no more than 10% > 400 per 100 ml	Water contact recreation	Criteria change or use clarification
Lost	Klamath Strait	0 to 9.8	Iron	Year Around	Table 20 Toxic Substances	Aquatic life; Human health	3
Lost	Klamath Strait	0 to 9.8	Manganese	Year Around	Table 20 Toxic Substances	Human health	3B
Lost	Klamath Strait	0 to 9.8	pH	Summer	pH 6.5 to 9.0	Anadromous fish passage; Resident fish and aquatic life; Salmonid fish rearing; Salmonid fish spawning; Water contact recreation	2
Lost	Klamath Strait	0 to 9.8	pH	FallWinterSpring	pH 6.5 to 9.0	Anadromous fish passage; Resident fish and aquatic life; Salmonid fish rearing; Salmonid fish spawning; Water contact recreation	2
Lost	Klamath Strait	0 to 9.8	Phosphate Phosphorus	Summer	Total phosphates as phosphorus <sup>4</sup>	Aquatic life	3B
Lost	Klamath Strait	0 to 9.8	Temperature	Summer	Previous narrative criteria: No measurable increase...	Anadromous fish passage; Resident fish and aquatic life; Salmonid fish rearing; Salmonid fish spawning	Criteria change or use clarification
Lost	Lost R. Diversion Canal	0 to 237.8	Dissolved Oxygen	Year Around	Cool water: Not less than 6.5 mg/l		5
Lost; Upper Klamath	Klamath R.	231.5 to 253	Ammonia	Year Around	Table 20 Toxic Substances	Aquatic life	5
Lost; Upper Klamath	Klamath R.	231.5 to 253	Chlorophyll a	Summer	Reservoir, river, estuary, non-thermally stratified lake: 0.015 mg/l	Aesthetics; Fishing; Livestock watering; Water contact recreation; Water supply	5
Lost; Upper Klamath	Klamath R.	231.1 to 251	Dissolved Oxygen	Year Around (Non-spawning)	Cool water: Not less than 6.5 mg/l	Cool-water aquatic life	5
Lost; Upper Klamath	Klamath R.	231 to 250	Fecal Coliform	Summer	Fecal coliform log mean of 200 organisms per 100 ml; no more than 10% > 400 per 100 ml	Water contact recreation	Attaining
Lost; Upper Klamath	Klamath R.	231 to 250	Fecal Coliform	FallWinterSpring	Fecal coliform log mean of 200 organisms per 100 ml; no more than 10% > 400 per 100 ml	Water contact recreation	Attaining
Lost; Upper	Klamath R.	231 to	Flow	Undefined	Creation of foul tastes, odors, or	Resident fish and aquatic life; Salmonid fish rearing;	4B

Watershed (USGS 4th Field Name)	Water Body (Stream/Lake)	River Miles	Parameter	Season	Criteria	Beneficial Uses	Integrated Category
Klamath		250	Modification		toxins. <sup>1</sup>	Salmonid fish spawning	
Lost; Upper Klamath	Klamath R.	231 to 250	Habitat Modification	Undefined	Creation of foul tastes, odors, or toxins. <sup>1</sup>	Resident fish and aquatic life; Salmonid fish rearing; 4B Salmonid fish spawning	
Lost; Upper Klamath	Klamath R.	231 to 250	Nutrients	Undefined		Aesthetics	3
Lost; Upper Klamath	Klamath R.	231 to 250	pH	FallWinterSpring	pH 6.5 to 9.0	Anadromous fish passage; Resident fish and aquatic life; Salmonid fish rearing; Salmonid fish spawning; Water contact recreation	Attaining
Lost; Upper Klamath	Klamath R.	231.5 to 253	pH	Summer	pH 6.5 to 9.0	Resident fish and aquatic life; Water contact recreation	5
Lost; Upper Klamath	Klamath R.	231 to 250	Sedimentation	Undefined	Bottom Deposits <sup>2</sup>	Resident fish and aquatic life; Salmonid fish rearing; 3 Salmonid fish spawning	3
Lost; Upper Klamath	Klamath R.	231 to 250	Temperature	Summer	Previous narrative criteria: No measurable increase...	Anadromous fish passage; Resident fish and aquatic life; Salmonid fish rearing; Salmonid fish spawning	Criteria change or use clarification
Lost; Upper Klamath	Klamath R.	231 to 250	Toxics	Undefined	Table 20 Toxic Substances	Anadromous fish passage; Drinking water; Resident fish and aquatic life	3
Lost; Upper Klamath	Klamath R. / Ewauna, Lake	232.7 to 253.7	Ammonia	Year Around	Table 20 Toxic Substances		5
Lost; Upper Klamath	Klamath R. / Ewauna, Lake	232.7 to 253.7	Dissolved Oxygen	Year Around (Non-spawning)	Cool water: Not less than 6.5 mg/l	Cool-water aquatic life	5
Lost; Upper Klamath	Klamath R. / Ewauna, Lake	232.7 to 253.7	pH	Summer	pH 6.0 to 8.5		5
Lost; Upper Klamath	Klamath R. / Ewauna, Lake	232.7 to 253.7	pH	FallWinterSpring	pH 6.0 to 8.5		5
Lost; Upper Klamath	Klamath R. / Ewauna, Lake	232.7 to 253.7	Phosphate Phosphorus	Summer	Total phosphates as phosphorus <sup>4</sup>	Aquatic life	3B
Sprague	Boulder Cr.	0 to 4.8	Temperature	Summer	Bull Trout: 10.0 C	Salmonid fish rearing; Salmonid fish spawning	4A
Sprague	Brownsworth Cr.	3.2 to 8.8	Temperature	Summer	Bull Trout: 10.0 C	Salmonid fish rearing; Salmonid fish spawning	4A
Sprague	Brownsworth Cr.	0 to 3.2	Temperature	Summer	Rearing: 17.8 C	Anadromous fish passage; Salmonid fish rearing	4A
Sprague	Buckboard Cr.	0 to 5	Temperature	Summer	Rearing: 17.8 C	Anadromous fish passage; Salmonid fish rearing	4A
Sprague	Calahan Cr.	0 to 7	Temperature	Summer	Rearing: 17.8 C	Anadromous fish passage; Salmonid fish rearing	4A
Sprague	Camp Cr.	0 to 3.1	Temperature	Summer	Rearing: 17.8 C	Anadromous fish passage; Salmonid fish rearing	3B
Sprague	Copperfield Cr.	0 to 3.2	Sedimentation	Summer	Bottom Deposits <sup>2</sup>	Resident fish and aquatic life; Salmonid fish rearing; 3 Salmonid fish spawning	3
Sprague	Corral Cr.	0 to 2.8	Temperature	Summer	Rearing: 17.8 C	Anadromous fish passage; Salmonid fish rearing	Attaining
Sprague	Coyote Cr.	0 to 10.4	Temperature	Summer	Rearing: 17.8 C	Anadromous fish passage; Salmonid fish rearing	4A

Watershed (USGS 4th Field Name)	Water Body (Stream/Lake)	River Miles	Parameter	Season	Criteria	Beneficial Uses	Integrated Category
Sprague	Deming Cr.	6.7 to 12.5	Temperature	Summer	Bull Trout: 10.0 C	Salmonid fish rearing; Salmonid fish spawning	4A
Sprague	Deming Cr.	0 to 6.7	Temperature	Summer	Rearing: 17.8 C	Anadromous fish passage; Salmonid fish rearing	4A
Sprague	Fishhole Cr.	0 to 25.6	Alkalinity	Year Around	Table 20 Toxic Substances	Aquatic life	3
Sprague	Fishhole Cr.	0 to 25.6	Ammonia	Year Around	Table 20 Toxic Substances	Aquatic life	3
Sprague	Fishhole Cr.	0 to 25.6	Biological Criteria	Year Around	Biocriteria <sup>5</sup>	Aquatic life	3B
Sprague	Fishhole Cr.	0 to 25.6	Chloride	Year Around	Table 20 Toxic Substances	Aquatic life	3
Sprague	Fishhole Cr.	0 to 25.6	Chlorophyll a	Summer	Reservoir, river, estuary, non-thermally stratified lake: 0.015 mg/l	Aesthetics; Fishing; Livestock watering; Water contact recreation; Water supply	3
Sprague	Fishhole Cr.	0 to 25.6	Dissolved Oxygen	Year Around (Non-spawning)	Cool water: Not less than 6.5 mg/l	Cool-water aquatic life	3
Sprague	Fishhole Cr.	0 to 25.6	E. Coli	Summer	E. Coli <sup>3</sup>	Water contact recreation	3
Sprague	Fishhole Cr.	0 to 25.6	Flow Modification	Undefined	Creation of foul tastes, odors, or toxins. <sup>1</sup>	Resident fish and aquatic life; Salmonid fish rearing; Salmonid fish spawning	4B
Sprague	Fishhole Cr.	0 to 25.6	Habitat Modification	Undefined	Creation of foul tastes, odors, or toxins. <sup>1</sup>	Resident fish and aquatic life; Salmonid fish rearing; Salmonid fish spawning	4B
Sprague	Fishhole Cr.	0 to 25.6	pH	Summer	pH 6.5 to 9.0	Resident fish and aquatic life; Water contact recreation	3
Sprague	Fishhole Cr.	0 to 25.6	Phosphate Phosphorus	Summer	Total phosphates as phosphorus <sup>4</sup>	Aquatic life	3
Sprague	Fishhole Cr.	0 to 25.6	Sedimentation	Undefined	Bottom Deposits <sup>2</sup>	Resident fish and aquatic life; Salmonid fish rearing; Salmonid fish spawning	3
Sprague	Fishhole Cr.	0 to 25.6	Temperature	Summer	Rearing: 17.8 C	Anadromous fish passage; Salmonid fish rearing	4A
Sprague	Fivemile Cr.	0 to 19.3	Flow Modification	Undefined	Creation of foul tastes, odors, or toxins. <sup>1</sup>	Resident fish and aquatic life; Salmonid fish rearing; Salmonid fish spawning	4B
Sprague	Fivemile Cr.	0 to 19.3	Habitat Modification	Undefined	Creation of foul tastes, odors, or toxins. <sup>1</sup>	Resident fish and aquatic life; Salmonid fish rearing; Salmonid fish spawning	4B
Sprague	Fivemile Cr.	0 to 19.3	Sedimentation	Undefined	Bottom Deposits <sup>2</sup>	Resident fish and aquatic life; Salmonid fish rearing; Salmonid fish spawning	3
Sprague	Fivemile Cr.	0 to 19.3	Temperature	Summer	Rearing: 17.8 C	Anadromous fish passage; Salmonid fish rearing	4A
Sprague	Leonard Cr.	0 to 3.1	Temperature	Summer	Bull Trout: 10.0 C	Salmonid fish rearing; Salmonid fish spawning	4A
Sprague	Long Cr.	0 to 15.6	Habitat Modification	Undefined	Creation of foul tastes, odors, or toxins. <sup>1</sup>	Resident fish and aquatic life; Salmonid fish rearing; Salmonid fish spawning	4B
Sprague	Long Cr.	0 to 15.6	Sedimentation	Undefined	Bottom Deposits <sup>2</sup>	Resident fish and aquatic life; Salmonid fish rearing; Salmonid fish spawning	3
Sprague	Long Cr.	0 to 15.6	Temperature	Summer	Rearing: 17.8 C	Anadromous fish passage; Salmonid fish rearing	4A

Watershed (USGS 4th Field Name)	Water Body (Stream/Lake)	River Miles	Parameter	Season	Criteria	Beneficial Uses	Integrated Category
Sprague	Meryl Cr.	0 to 15	Flow Modification	Undefined	Creation of foul tastes, odors, or toxins. <sup>1</sup>	Resident fish and aquatic life; Salmonid fish rearing; 4B Salmonid fish spawning	
Sprague	Meryl Cr.	0 to 15	Sedimentation	Undefined	Bottom Deposits <sup>2</sup>	Resident fish and aquatic life; Salmonid fish rearing; 3 Salmonid fish spawning	
Sprague	Meryl Cr.	0 to 15	Temperature	Undefined	Rearing: 17.8 C	Anadromous fish passage; Salmonid fish rearing	3
Sprague	N.F. Sprague R.	0 to 18	Alkalinity	Year Around	Table 20 Toxic Substances	Aquatic life	2
Sprague	N.F. Sprague R.	18 to 33.5	Alkalinity	Year Around	Table 20 Toxic Substances	Aquatic life	3B
Sprague	N.F. Sprague R.	0 to 33.5	Ammonia	Year Around	Table 20 Toxic Substances	Aquatic life	2
Sprague	N.F. Sprague R.	0 to 33.5	Chloride	Year Around	Table 20 Toxic Substances	Aquatic life	3
Sprague	N.F. Sprague R.	0 to 33.5	Chlorophyll a	FallWinterSpring	Reservoir, river, estuary, non-thermally stratified lake: 0.015 mg/l	Aesthetics; Fishing; Livestock watering; Water contact recreation; Water supply	3
Sprague	N.F. Sprague R.	0 to 33.5	Chlorophyll a	Summer	Reservoir, river, estuary, non-thermally stratified lake: 0.015 mg/l	Aesthetics; Fishing; Livestock watering; Water contact recreation; Water supply	3
Sprague	N.F. Sprague R.	0 to 8	Dissolved Oxygen	Year Around (Non-spawning)	Cool water: Not less than 6.5 mg/l	Cool-water aquatic life	2
Sprague	N.F. Sprague R.	8 to 11.3	Dissolved Oxygen	Year Around (Non-spawning)	Cold water: Not less than 8.0 mg/l or 90% of saturation	Cold-water aquatic life	3
Sprague	N.F. Sprague R.	0 to 33.5	Dissolved Oxygen	January 1 - May 15	Spawning: Not less than 11.0 mg/L or 95% of saturation	Resident trout spawning	3
Sprague	N.F. Sprague R.	0 to 33.5	E. Coli	Summer	30-day log mean of 126 E. coli organisms per 100 ml; no single sample > 406 organisms per 100 ml	Water contact recreation	2
Sprague	N.F. Sprague R.	0 to 33.5	E. Coli	FallWinterSpring	E. Coli <sup>3</sup>	Water contact recreation	3
Sprague	N.F. Sprague R.	0 to 33.5	Flow Modification	Undefined	Creation of foul tastes, odors, or toxins. <sup>1</sup>	Resident fish and aquatic life; Salmonid fish rearing; 4B Salmonid fish spawning	
Sprague	N.F. Sprague R.	0 to 33.5	Habitat Modification	Undefined	Creation of foul tastes, odors, or toxins. <sup>1</sup>	Resident fish and aquatic life; Salmonid fish rearing; 4B Salmonid fish spawning	
Sprague	N.F. Sprague R.	0 to 33.5	Iron	Year Around	Table 20 Toxic Substances	Aquatic life; Human health	3
Sprague	N.F. Sprague R.	0 to 33.5	Manganese	Year Around	Table 20 Toxic Substances	Human health	3
Sprague	N.F. Sprague R.	0 to 33.5	pH	Summer	pH 6.5 to 9.0	Resident fish and aquatic life; Water contact recreation	2
Sprague	N.F. Sprague R.	0 to 33.5	pH	FallWinterSpring	pH 6.5 to 9.0	Resident fish and aquatic life; Water contact recreation	3
Sprague	N.F. Sprague R.	0 to 33.5	Phosphate Phosphorus	Summer	Total phosphates as phosphorus <sup>4</sup>	Aquatic life	3B
Sprague	N.F. Sprague R.	0 to 33.5	Sedimentation	Undefined	Bottom Deposits <sup>2</sup>	Resident fish and aquatic life; Salmonid fish rearing; 3 Salmonid fish spawning	



Watershed (USGS 4th Field Name)	Water Body (Stream/Lake)	River Miles	Parameter	Season	Criteria	Beneficial Uses	Integrated Category
Sprague	N.F. Sprague R.	28.3 to 33.5	Temperature	Summer	Rearing: 17.8 C	Anadromous fish passage; Salmonid fish rearing	Attaining
Sprague	N.F. Sprague R.	0 to 33.5	Temperature	Summer	Rearing: 17.8 C	Anadromous fish passage; Salmonid fish rearing	4A
Sprague	Paradise Cr.	0 to 8.6	Temperature	Summer	Rearing: 17.8 C	Anadromous fish passage; Salmonid fish rearing	4A
Sprague	Pothole Cr.	0 to 6.1	Temperature	Summer	Rearing: 17.8 C	Anadromous fish passage; Salmonid fish rearing	4A
Sprague	S.F. Sprague R.	0 to 31.3	Alkalinity	Year Around	Table 20 Toxic Substances	Aquatic life	2
Sprague	S.F. Sprague R.	0 to 31.3	Ammonia	Year Around	Table 20 Toxic Substances	Aquatic life	2
Sprague	S.F. Sprague R.	0 to 27.7	Aquatic Weeds Or Algae	Undefined	The development of fungi or other growths <sup>6</sup>	Aesthetics; Fishing; Water contact recreation	3
Sprague	S.F. Sprague R.	0 to 31.3	Chloride	Year Around	Table 20 Toxic Substances	Aquatic life	3
Sprague	S.F. Sprague R.	0 to 31.3	Chlorophyll a	FallWinterSpring	Reservoir, river, estuary, non-thermally stratified lake: 0.015 mg/l	Aesthetics; Fishing; Livestock watering; Water contact recreation; Water supply	3
Sprague	S.F. Sprague R.	0 to 31.3	Chlorophyll a	Summer	Reservoir, river, estuary, non-thermally stratified lake: 0.015 mg/l	Aesthetics; Fishing; Livestock watering; Water contact recreation; Water supply	3
Sprague	S.F. Sprague R.	0 to 14.3	Dissolved Oxygen	Year Around (Non-spawning)	Cool water: Not less than 6.5 mg/l	Cool-water aquatic life	3
Sprague	S.F. Sprague R.	0 to 31.2	Dissolved Oxygen	January 1 - May 15	Spawning: Not less than 11.0 mg/L or 95% of saturation	Resident trout spawning	3
Sprague	S.F. Sprague R.	0 to 31.3	E. Coli	FallWinterSpring	E. Coli <sup>3</sup>	Water contact recreation	3
Sprague	S.F. Sprague R.	0 to 31.3	E. Coli	Summer	E. Coli <sup>3</sup>	Water contact recreation	5
Sprague	S.F. Sprague R.	0 to 27.7	Flow Modification	Undefined	Creation of foul tastes, odors, or toxins. <sup>1</sup>	Resident fish and aquatic life; Salmonid fish rearing; Salmonid fish spawning	4B
Sprague	S.F. Sprague R.	0 to 27.7	Habitat Modification	Undefined	Creation of foul tastes, odors, or toxins. <sup>1</sup>	Resident fish and aquatic life; Salmonid fish rearing; Salmonid fish spawning	4B
Sprague	S.F. Sprague R.	0 to 31.3	Iron	Year Around	Table 20 Toxic Substances	Aquatic life; Human health	3
Sprague	S.F. Sprague R.	0 to 31.3	Manganese	Year Around	Table 20 Toxic Substances	Human health	3
Sprague	S.F. Sprague R.	0 to 27.7	Nutrients	Undefined		Aesthetics	3
Sprague	S.F. Sprague R.	0 to 31.3	pH	Summer	pH 6.5 to 9.0	Resident fish and aquatic life; Water contact recreation	3
Sprague	S.F. Sprague R.	0 to 31.3	pH	FallWinterSpring	pH 6.5 to 9.0	Resident fish and aquatic life; Water contact recreation	3
Sprague	S.F. Sprague R.	0 to 31.2	Phosphate Phosphorus	Summer	Total phosphates as phosphorus <sup>4</sup>	Aquatic life	3B
Sprague	S.F. Sprague R.	0 to 27.7	Sedimentation	Undefined	Bottom Deposits <sup>2</sup>	Resident fish and aquatic life; Salmonid fish rearing; Salmonid fish spawning	3
Sprague	S.F. Sprague R.	27.7 to	Temperature	Summer	Rearing: 17.8 C	Anadromous fish passage; Salmonid fish rearing	Attaining

Watershed (USGS 4th Field Name)	Water Body (Stream/Lake)	River Miles	Parameter	Season	Criteria	Beneficial Uses	Integrated Category
		31.2					
Sprague	S.F. Sprague R.	0 to 27.7	Temperature	Summer	Rearing: 17.8 C	Anadromous fish passage; Salmonid fish rearing	4A
Sprague	Sprague R.	0 to 79.2	Alkalinity	Year Around	Table 20 Toxic Substances	Aquatic life	2
Sprague	Sprague R.	0 to 79.2	Ammonia	Year Around	Table 20 Toxic Substances	Aquatic life	2
Sprague	Sprague R.	0 to 79.2	Aquatic Weeds Or Algae	Undefined	The development of fungi or other growths <sup>6</sup>	Aesthetics; Fishing; Water contact recreation	3
Sprague	Sprague R.	0 to 79.2	Chloride	Year Around	Table 20 Toxic Substances	Aquatic life	3
Sprague	Sprague R.	0 to 79.2	Chlorophyll a	FallWinterSpring	Reservoir, river, estuary, non-thermally stratified lake: 0.015 mg/l	Aesthetics; Fishing; Livestock watering; Water contact recreation; Water supply	3
Sprague	Sprague R.	0 to 79.2	Chlorophyll a	Summer	Reservoir, river, estuary, non-thermally stratified lake: 0.015 mg/l	Aesthetics; Fishing; Livestock watering; Water contact recreation; Water supply	3
Sprague	Sprague R.	45.7 to 79.2	Dissolved Oxygen	Year Around (Non-spawning)	Cool water: Not less than 6.5 mg/l	Cool-water aquatic life	2
Sprague	Sprague R.	0 to 45.7	Dissolved Oxygen	Year Around (Non-spawning)	Cold water: Not less than 8.0 mg/l or 90% of saturation	Cold-water aquatic life	4A
Sprague	Sprague R.	0 to 79.2	Dissolved Oxygen	Summer	Cold water: Not less than 8.0 mg/l or 90% of saturation	Anadromous fish passage; Salmonid fish rearing	4A
Sprague	Sprague R.	0 to 79.2	Dissolved Oxygen	January 1 - May 15	Spawning: Not less than 11.0 mg/L or 95% of saturation	Resident trout spawning	3
Sprague	Sprague R.	0 to 79.2	E. Coli	Summer	30-day log mean of 126 E. coli organisms per 100 ml; no single sample > 406 organisms per 100 ml	Water contact recreation	2
Sprague	Sprague R.	0 to 79.2	E. Coli	FallWinterSpring	E. Coli <sup>3</sup>	Water contact recreation	3
Sprague	Sprague R.	0 to 79.2	Flow Modification	Undefined	Creation of foul tastes, odors, or toxins. <sup>1</sup>	Resident fish and aquatic life; Salmonid fish rearing; Salmonid fish spawning	4B
Sprague	Sprague R.	0 to 79.2	Habitat Modification	Undefined	Creation of foul tastes, odors, or toxins. <sup>1</sup>	Resident fish and aquatic life; Salmonid fish rearing; Salmonid fish spawning	4B
Sprague	Sprague R.	0 to 79.2	Iron	Year Around	Table 20 Toxic Substances	Aquatic life; Human health	3
Sprague	Sprague R.	0 to 79.2	Manganese	Year Around	Table 20 Toxic Substances	Human health	3
Sprague	Sprague R.	0 to 79.2	Nutrients	Undefined		Aesthetics	3
Sprague	Sprague R.	0 to 79.2	pH	FallWinterSpring	pH 6.5 to 9.0	Resident fish and aquatic life; Water contact recreation	3
Sprague	Sprague R.	0 to 79.2	pH	Summer	pH 6.5 to 9.0	Anadromous fish passage; Resident fish and aquatic life; Salmonid fish rearing; Salmonid fish spawning; Water contact recreation	4A
Sprague	Sprague R.	0 to 79.2	Phosphate Phosphorus	Summer	Total phosphates as phosphorus <sup>4</sup>	Aquatic life	3B

Watershed (USGS 4th Field Name)	Water Body (Stream/Lake)	River Miles	Parameter	Season	Criteria	Beneficial Uses	Integrated Category
Sprague	Sprague R.	0 to 79.2	Sedimentation	Undefined	Bottom Deposits <sup>2</sup>	Resident fish and aquatic life; Salmonid fish rearing; 3 Salmonid fish spawning	
Sprague	Sprague R.	0 to 79.2	Temperature	Summer	Rearing: 17.8 C	Anadromous fish passage; Salmonid fish rearing	4A
Sprague	Sycan R.	0 to 71.4	Alkalinity	Year Around	Table 20 Toxic Substances	Aquatic life	2
Sprague	Sycan R.	0 to 71.4	Ammonia	Year Around	Table 20 Toxic Substances	Aquatic life	2
Sprague	Sycan R.	0 to 64.1	Aquatic Weeds Or Algae	Undefined	The development of fungi or other growths <sup>6</sup>	Aesthetics; Fishing; Water contact recreation	3
Sprague	Sycan R.	0 to 71.4	Chloride	Year Around	Table 20 Toxic Substances	Aquatic life	3
Sprague	Sycan R.	0 to 71.4	Chlorophyll a	Summer	Reservoir, river, estuary, non-thermally stratified lake: 0.015 mg/l	Aesthetics; Fishing; Livestock watering; Water contact recreation; Water supply	3
Sprague	Sycan R.	0 to 71.4	Chlorophyll a	FallWinterSpring	Reservoir, river, estuary, non-thermally stratified lake: 0.015 mg/l	Aesthetics; Fishing; Livestock watering; Water contact recreation; Water supply	3
Sprague	Sycan R.	0 to 64.1	Dissolved Oxygen	Year Around (Non-spawning)	Cool water: Not less than 6.5 mg/l	Cool-water aquatic life	2
Sprague	Sycan R.	0 to 71.4	Dissolved Oxygen	January 1 - May 15	Spawning: Not less than 11.0 mg/L or 95% of saturation	Resident trout spawning	3
Sprague	Sycan R.	0 to 71.4	E. Coli	Summer	30-day log mean of 126 E. coli organisms per 100 ml; no single sample > 406 organisms per 100 ml	Water contact recreation	2
Sprague	Sycan R.	0 to 71.4	E. Coli	FallWinterSpring	E. Coli <sup>3</sup>	Water contact recreation	3
Sprague	Sycan R.	0 to 64.1	Flow Modification	Undefined	Creation of foul tastes, odors, or toxins. <sup>1</sup>	Resident fish and aquatic life; Salmonid fish rearing; 4B Salmonid fish spawning	
Sprague	Sycan R.	0 to 64.1	Habitat Modification	Undefined	Creation of foul tastes, odors, or toxins. <sup>1</sup>	Resident fish and aquatic life; Salmonid fish rearing; 4B Salmonid fish spawning	
Sprague	Sycan R.	0 to 71.4	Iron	Year Around	Table 20 Toxic Substances	Aquatic life; Human health	3
Sprague	Sycan R.	0 to 71.4	Manganese	Year Around	Table 20 Toxic Substances	Human health	3
Sprague	Sycan R.	0 to 64.1	Nutrients	Undefined		Aesthetics	3
Sprague	Sycan R.	0 to 71.4	pH	Summer	pH 6.5 to 9.0	Resident fish and aquatic life; Water contact recreation	2
Sprague	Sycan R.	0 to 71.4	pH	FallWinterSpring	pH 6.5 to 9.0	Resident fish and aquatic life; Water contact recreation	3
Sprague	Sycan R.	0 to 71.4	Phosphate Phosphorus	Summer	Total phosphates as phosphorus <sup>4</sup>	Aquatic life	3B
Sprague	Sycan R.	0 to 64.1	Sedimentation	Undefined	Bottom Deposits <sup>2</sup>	Resident fish and aquatic life; Salmonid fish rearing; 3 Salmonid fish spawning	
Sprague	Sycan R.	0 to 64.1	Temperature	Summer	Rearing: 17.8 C	Anadromous fish passage; Salmonid fish rearing	4A
Sprague	Trout Cr.	0 to 1.4	Ammonia	Year Around	Table 20 Toxic Substances	Aquatic life	3

Watershed (USGS 4th Field Name)	Water Body (Stream/Lake)	River Miles	Parameter	Season	Criteria	Beneficial Uses	Integrated Category
Sprague	Trout Cr.	0 to 1.4	Dissolved Oxygen	Year Around (Non-spawning)	Cold water: Not less than 8.0 mg/l or 90% of saturation	Cold-water aquatic life	3
Sprague	Trout Cr.	0 to 1.4	E. Coli	Summer	E. Coli <sup>3</sup>	Water contact recreation	3
Sprague	Trout Cr.	0 to 1.4	pH	Summer	pH 6.5 to 9.0	Resident fish and aquatic life; Water contact recreation	3
Sprague	Trout Cr.	0 to 1.4	Phosphate Phosphorus	Summer	Total phosphates as phosphorus <sup>4</sup>	Aquatic life	3
Sprague	Trout Cr.	0 to 1.4	Temperature	Summer	Rearing: 17.8 C	Anadromous fish passage; Salmonid fish rearing	4A
Upper Klamath	Beaver Cr.	0 to 5.5	Temperature	Year Around (Non-spawning)	Redband or Lahontan cutthroat trout: 20.0 degrees Celsius 7-day-average maximum	Redband or Lahontan cutthroat trout	5
Upper Klamath	Clover Cr.	0 to 8.4	Biological Criteria	Year Around	Biocriteria <sup>5</sup>	Resident fish and aquatic life	3
Upper Klamath	Clover Cr.	0 to 8.4	Habitat Modification	Undefined	Creation of foul tastes, odors, or toxins. <sup>1</sup>	Resident fish and aquatic life; Salmonid fish rearing; 4B Salmonid fish spawning	
Upper Klamath	Clover Cr.	0 to 8.4	Sedimentation	Undefined	Bottom Deposits <sup>2</sup>	Resident fish and aquatic life; Salmonid fish rearing; 303(d) Salmonid fish spawning	
Upper Klamath	Clover Cr.	0 to 8.4	Temperature	Undefined		Anadromous fish passage; Resident fish and aquatic life; Salmonid fish rearing; Salmonid fish spawning	3
Upper Klamath	Corral Cr.	0 to 2.4	Temperature	Summer	Rearing: 17.8 C	Anadromous fish passage; Salmonid fish rearing	3B
Upper Klamath	Grizzly Cr.	0 to 3	Temperature	Summer	Rearing: 17.8 C	Anadromous fish passage; Salmonid fish rearing	303(d)
Upper Klamath	Hoxie Cr.	0.8 to 4.4	Temperature	Summer	Rearing: 17.8 C	Anadromous fish passage; Salmonid fish rearing	303(d)
Upper Klamath	Jenny Cr.	0 to 17.8	Sedimentation	Undefined	Bottom Deposits <sup>2</sup>	Resident fish and aquatic life; Salmonid fish rearing; 3 Salmonid fish spawning	
Upper Klamath	Jenny Cr.	0 to 18.7	Temperature	Summer	Rearing: 17.8 C	Anadromous fish passage; Salmonid fish rearing	Attaining
Upper Klamath	Jenny Cr.	0 to 17.8	Temperature	Summer	Rearing: 17.8 C	Anadromous fish passage; Salmonid fish rearing	303(d)
Upper Klamath	Johnson Cr.	0 to 9.4	Temperature	Summer	Rearing: 17.8 C	Anadromous fish passage; Salmonid fish rearing	303(d)
Upper Klamath	Keene Cr.	0 to 7.2	Temperature	Summer	Rearing: 17.8 C	Anadromous fish passage; Salmonid fish rearing	303(d)
Upper Klamath	Keene Cr.	7.5 to 9.7	Temperature	Summer	Rearing: 17.8 C	Anadromous fish passage; Salmonid fish rearing	303(d)
Upper Klamath	Keene Cr. / Hyatt Reservoir	11.3896 to 13.8	Aquatic Weeds Or Algae	Undefined	The development of fungi or other growths <sup>6</sup>	Aesthetics; Drinking water; Fishing; Livestock watering; Water contact recreation; Water supply	5
Upper Klamath	Keene Cr. / Hyatt Reservoir	11.3896 to 13.8	Nutrients	Undefined		Aesthetics	3
Upper Klamath	Klamath R.	207 to 231.5	Ammonia	Year Around	Table 20 Toxic Substances	Aquatic life	2
Upper Klamath	Klamath R.	207 to 231	Chlorophyll a	Summer	Reservoir, river, estuary, non-thermally stratified lake: 0.015 mg/l	Aesthetics; Fishing; Livestock watering; Water contact recreation; Water supply	Attaining

<b>Watershed (USGS 4th Field Name)</b>	<b>Water Body (Stream/Lake)</b>	<b>River Miles</b>	<b>Parameter</b>	<b>Season</b>	<b>Criteria</b>	<b>Beneficial Uses</b>	<b>Integrated Category</b>
Upper Klamath	Klamath R.	207 to 231.1	Dissolved Oxygen	Year Around (Non-spawning)	Cold water: Not less than 8.0 mg/l or 90% of saturation	Cold-water aquatic life	5
Upper Klamath	Klamath R.	207 to 231.1	Dissolved Oxygen	January 1 - May 15	Spawning: Not less than 11.0 mg/L or 95% of saturation	Resident trout spawning	5
Upper Klamath	Klamath R.	207 to 231	Fecal Coliform	Summer	Fecal coliform log mean of 200 organisms per 100 ml; no more than 10% > 400 per 100 ml	Water contact recreation	Attaining
Upper Klamath	Klamath R.	207 to 231	Fecal Coliform	FallWinterSpring	Fecal coliform log mean of 200 organisms per 100 ml; no more than 10% > 400 per 100 ml	Water contact recreation	Attaining
Upper Klamath	Klamath R.	207 to 231	Habitat Modification	Undefined	Creation of foul tastes, odors, or toxins. <sup>1</sup>	Resident fish and aquatic life; Salmonid fish rearing; 4B Salmonid fish spawning	
Upper Klamath	Klamath R.	207 to 231	Nutrients	Undefined		Aesthetics	3
Upper Klamath	Klamath R.	207 to 231	pH	Summer	pH 6.5 to 9.0	Resident fish and aquatic life; Anadromous fish passage; Water contact recreation; Salmonid fish spawning; Salmonid fish rearing	Attaining
Upper Klamath	Klamath R.	207 to 231	pH	FallWinterSpring	pH 6.5 to 9.0	Salmonid fish rearing; Salmonid fish spawning; Resident fish and aquatic life; Anadromous fish passage; Water contact recreation	Attaining
Upper Klamath	Klamath R.	207 to 231.5	pH	Summer	pH 6.5 to 9.0	Resident fish and aquatic life; Water contact recreation	2
Upper Klamath	Klamath R.	207 to 231	Sedimentation	Undefined	Bottom Deposits <sup>2</sup>	Resident fish and aquatic life; Salmonid fish rearing; 3 Salmonid fish spawning	
Upper Klamath	Klamath R.	207 to 231.1	Temperature	Year Around (Non-spawning)	Redband or Lahontan cutthroat trout: 20.0 degrees Celsius 7-day- average maximum	Redband or Lahontan cutthroat trout	5
Upper Klamath	Klamath R. / Unnamed Lake (J.C. Boyle Reservoir)	224.7 to 228.1	Chlorophyll a	Summer	Reservoir, river, estuary, non- thermally stratified lake: 0.015 mg/l	Aesthetics; Fishing; Livestock watering; Water contact recreation; Water supply	3
Upper Klamath	Klamath R. / Unnamed Lake (J.C. Boyle Reservoir)	224.7 to 228.1	Nutrients	Undefined		Aesthetics	3
Upper Klamath	Klamath R. / Unnamed Lake (J.C. Boyle Reservoir)	224.7 to 228.1	pH	Summer	pH 6.0 to 8.5	Resident fish and aquatic life; Water contact recreation	3

<b>Watershed (USGS 4th Field Name)</b>	<b>Water Body (Stream/Lake)</b>	<b>River Miles</b>	<b>Parameter</b>	<b>Season</b>	<b>Criteria</b>	<b>Beneficial Uses</b>	<b>Integrated Category</b>
Upper Klamath	Lincoln Cr.	0 to 2.7	Temperature	Summer	Rearing: 17.8 C	Anadromous fish passage; Salmonid fish rearing	3B
Upper Klamath	Long Prairie Cr.	0 to 11.9	Sedimentation	Undefined	Bottom Deposits <sup>2</sup>	Resident fish and aquatic life; Salmonid fish rearing; 3 Salmonid fish spawning	
Upper Klamath	Mill Cr.	0 to 3.9	Temperature	Summer	Rearing: 17.8 C	Anadromous fish passage; Salmonid fish rearing	303(d)
Upper Klamath	Miners Cr.	0 to 4.3	Sedimentation	Undefined	Bottom Deposits <sup>2</sup>	Resident fish and aquatic life; Salmonid fish rearing; 303(d) Salmonid fish spawning	
Upper Klamath	Miners Cr.	0 to 4.3	Temperature	Year Around (Non-spawning)	Redband or Lahontan cutthroat trout: 20.0 degrees Celsius 7-day- average maximum	Redband or Lahontan cutthroat trout	2
Upper Klamath	S.F. Keene Cr.	0 to 3.1	Temperature	Summer	Rearing: 17.8 C	Anadromous fish passage; Salmonid fish rearing	303(d)
Upper Klamath	Spencer Cr.	0 to 18.9	Biological Criteria	Year Around	Biocriteria <sup>5</sup>	Resident fish and aquatic life	303(d)
Upper Klamath	Spencer Cr.	0 to 18.9	Flow Modification	Undefined	Creation of foul tastes, odors, or toxins. <sup>1</sup>	Resident fish and aquatic life; Salmonid fish rearing; 4B Salmonid fish spawning	
Upper Klamath	Spencer Cr.	0 to 18.9	Habitat Modification	Undefined	Creation of foul tastes, odors, or toxins. <sup>1</sup>	Resident fish and aquatic life; Salmonid fish rearing; 4B Salmonid fish spawning	
Upper Klamath	Spencer Cr.	0 to 18.9	Sedimentation	Undefined	Bottom Deposits <sup>2</sup>	Resident fish and aquatic life; Salmonid fish rearing; 303(d) Salmonid fish spawning	
Upper Klamath	Spencer Cr.	0 to 18.9	Temperature	Year Around (Non-spawning)	Redband or Lahontan cutthroat trout: 20.0 degrees Celsius 7-day- average maximum	Redband or Lahontan cutthroat trout	5
UKL	Annie Cr.	0 to 16.3	Biological Criteria	Year Around	Biocriteria <sup>5</sup>	Aquatic life	5
UKL	Annie Cr.	0 to 6.1	Flow Modification	Undefined	Creation of foul tastes, odors, or toxins. <sup>1</sup>	Resident fish and aquatic life; Salmonid fish rearing; 4B Salmonid fish spawning	
UKL	Annie Cr.	0 to 6.1	Habitat Modification	Undefined	Creation of foul tastes, odors, or toxins. <sup>1</sup>	Resident fish and aquatic life; Salmonid fish rearing; 4B Salmonid fish spawning	
UKL	Annie Cr.	0 to 6.1	Temperature	Undefined		Anadromous fish passage; Resident fish and aquatic life; Salmonid fish rearing; Salmonid fish spawning	3
UKL	Cherry Cr.	0 to 9.7	Flow Modification	Undefined	Creation of foul tastes, odors, or toxins. <sup>1</sup>	Resident fish and aquatic life; Salmonid fish rearing; 4B Salmonid fish spawning	
UKL	Cherry Cr.	0 to 9.7	Temperature	Summer	Rearing: 17.8 C	Anadromous fish passage; Salmonid fish rearing	Attaining
UKL	Crooked Cr.	0 to 9	pH	Summer	pH 6.5 to 9.0	Resident fish and aquatic life; Water contact recreation	3
UKL	Crooked Cr.	0 to 9	Phosphate Phosphorus	Summer	Total phosphates as phosphorus <sup>4</sup>	Aquatic life	3
UKL	Fourmile Cr.	0 to 10.2	Flow Modification	Undefined	Creation of foul tastes, odors, or toxins. <sup>1</sup>	Resident fish and aquatic life; Salmonid fish rearing; 4B Salmonid fish spawning	

Watershed (USGS 4th Field Name)	Water Body (Stream/Lake)	River Miles	Parameter	Season	Criteria	Beneficial Uses	Integrated Category
UKL	Fourmile Cr.	1 to 10.2	Temperature	Summer	Rearing: 17.8 C	Anadromous fish passage; Salmonid fish rearing	Attaining
UKL	Fourmile Cr.	0 to 1	Temperature	Summer	Rearing: 17.8 C	Anadromous fish passage; Salmonid fish rearing	4A
UKL	Klamath R. / Agency Lake	275 to 282	Aquatic Weeds Or Algae	Undefined	The development of fungi or other growths <sup>6</sup>	Aesthetics; Drinking water; Fishing; Livestock watering; Water contact recreation; Water supply	4A
UKL	Klamath R. / Agency Lake	275 to 282	Chlorophyll a	Summer	Thermally stratified lake: 0.01 mg/l	Aesthetics; Fishing; Livestock watering; Water contact recreation; Water supply	4A
UKL	Klamath R. / Agency Lake	275 to 282	Dissolved Oxygen	Summer	Cool water: Not less than 6.5 mg/l	Resident fish and aquatic life; Salmonid fish rearing	4A
UKL	Klamath R. / Agency Lake	275 to 282	Nutrients	Undefined		Aesthetics	3
UKL	Klamath R. / Agency Lake	275 to 282	pH	Summer	pH 6.5 to 8.5	Anadromous fish passage; Resident fish and aquatic life; Salmonid fish rearing; Salmonid fish spawning; Water contact recreation	4A
UKL	Klamath R. / Agency Lake	275 to 282	Sedimentation	Undefined	Bottom Deposits <sup>2</sup>	Resident fish and aquatic life; Salmonid fish rearing; Salmonid fish spawning	3
UKL	Klamath R. / UKL	254.9 to 278.5	Aquatic Weeds Or Algae	Undefined	The development of fungi or other growths <sup>6</sup>	Aesthetics; Drinking water; Fishing; Livestock watering; Water contact recreation; Water supply	4A
UKL	Klamath R. / UKL	254.9 to 278.5	Chlorophyll a	Summer	Thermally stratified lake: 0.01 mg/l	Aesthetics; Fishing; Livestock watering; Water contact recreation; Water supply	4A
UKL	Klamath R. / UKL	254.9 to 278.5	Dissolved Oxygen	Summer	Cool water: Not less than 6.5 mg/l	Resident fish and aquatic life; Salmonid fish rearing; Salmonid fish spawning	4A
UKL	Klamath R. / UKL	254.9 to 278.5	Nutrients	Undefined		Aesthetics	3
UKL	Klamath R. / UKL	254.9 to 278.5	pH	Summer	pH 6.5 to 8.5	Anadromous fish passage; Resident fish and aquatic life; Salmonid fish rearing; Salmonid fish spawning; Water contact recreation	4A
UKL	Klamath R. / UKL	254.9 to 278.5	Sedimentation	Undefined	Bottom Deposits <sup>2</sup>	Resident fish and aquatic life; Salmonid fish rearing; Salmonid fish spawning	3
UKL	Klamath R. / UKL	254.9 to 278.5	Temperature	Undefined		Anadromous fish passage; Resident fish and aquatic life; Salmonid fish rearing; Salmonid fish spawning	3
UKL	Rock Cr.	0 to 5.7	Biological Criteria	Year Around	Biocriteria: Waters of the state must be of sufficient quality to support aquatic species without detrimental changes in the resident biological communities.	Aquatic life	2
UKL	Rock Cr.	0 to 5.7	Habitat Modification	Undefined	Creation of foul tastes, odors, or toxins. <sup>1</sup>	Resident fish and aquatic life; Salmonid fish rearing; Salmonid fish spawning	4B
UKL	Rock Cr.	0 to 5.7	Temperature	Summer	Rearing: 17.8 C	Anadromous fish passage; Salmonid fish rearing	4A

Watershed (USGS 4th Field Name)	Water Body (Stream/Lake)	River Miles	Parameter	Season	Criteria	Beneficial Uses	Integrated Category
UKL	Sevenmile Canal	0 to 10.5	Dissolved Oxygen	Year Around	Cool water: Not less than 6.5 mg/l		5
UKL	Sevenmile Cr.	4.2 to 12.7	Temperature	Summer	Rearing: 17.8 C	Anadromous fish passage; Salmonid fish rearing	Attaining
UKL	Sevenmile Ditch	0 to 1.8	Dissolved Oxygen	Undefined		Anadromous fish passage; Resident fish and aquatic life; Salmonid fish rearing; Salmonid fish spawning	3
UKL	Sevenmile Ditch	0 to 1.8	Flow Modification	Undefined	Creation of foul tastes, odors, or toxins. <sup>1</sup>	Resident fish and aquatic life; Salmonid fish rearing; 4B Salmonid fish spawning	
UKL	Sevenmile Ditch	0 to 1.8	Habitat Modification	Undefined	Creation of foul tastes, odors, or toxins. <sup>1</sup>	Resident fish and aquatic life; Salmonid fish rearing; 4B Salmonid fish spawning	
UKL	Sevenmile Ditch	0 to 1.8	Nutrients	Undefined		Aesthetics	3
UKL	Sevenmile Ditch	0 to 1.8	Sedimentation	Undefined	Bottom Deposits <sup>2</sup>	Resident fish and aquatic life; Salmonid fish rearing; 3 Salmonid fish spawning	
UKL	Sevenmile Ditch	0 to 1.8	Temperature	Undefined		Anadromous fish passage; Resident fish and aquatic life; Salmonid fish rearing; Salmonid fish spawning	3
UKL	Sun Cr.	0 to 13.6	Temperature	Undefined		Anadromous fish passage; Resident fish and aquatic life; Salmonid fish rearing; Salmonid fish spawning	3
UKL	Threemile Cr.	0 to 7.6	Habitat Modification	Undefined	Creation of foul tastes, odors, or toxins. <sup>1</sup>	Resident fish and aquatic life; Salmonid fish rearing; 4B Salmonid fish spawning	
UKL	Threemile Cr.	0 to 7.6	Temperature	Summer	Rearing: 17.8 C	Anadromous fish passage; Salmonid fish rearing	Attaining
UKL	Wood R.	0 to 17.9	Alkalinity	Year Around	Table 20 Toxic Substances	Aquatic life	3
UKL	Wood R.	0 to 17.9	Ammonia	Year Around	Table 20 Toxic Substances	Aquatic life	2
UKL	Wood R.	0 to 17.8	Chlorophyll a	Summer	Thermally stratified lake: 0.01 mg/l	Aesthetics; Fishing; Livestock watering; Water contact recreation; Water supply	Attaining
UKL	Wood R.	0 to 15	Dissolved Oxygen	Year Around (Non-spawning)	Cool water: Not less than 6.5 mg/l	Cool-water aquatic life	3
UKL	Wood R.	15 to 17.8	Dissolved Oxygen	Year Around (Non-spawning)	Cool water: Not less than 6.5 mg/l	Cool-water aquatic life	3
UKL	Wood R.	0 to 17.8	Dissolved Oxygen	August 15 - June 15	Spawning: Not less than 11.0 mg/L or 95% of saturation	Resident trout spawning; Bull trout spawning and juvenile rearing	3
UKL	Wood R.	0 to 17.8	Flow Modification	Undefined	Creation of foul tastes, odors, or toxins. <sup>1</sup>	Resident fish and aquatic life; Salmonid fish rearing; 4B Salmonid fish spawning	
UKL	Wood R.	0 to 17.8	Habitat Modification	Undefined	Creation of foul tastes, odors, or toxins. <sup>1</sup>	Resident fish and aquatic life; Salmonid fish rearing; 4B Salmonid fish spawning	
UKL	Wood R.	0 to 17.8	Nutrients	Undefined		Aesthetics	3B
UKL	Wood R.	0 to 17.8	pH	Undefined	pH 6.5 to 9.0	Anadromous fish passage; Resident fish and aquatic life; Salmonid fish rearing; Salmonid fish spawning;	Attaining



Watershed (USGS 4th Field Name)	Water Body (Stream/Lake)	River Miles	Parameter	Season	Criteria	Beneficial Uses	Integrated Category
						Water contact recreation	
UKL	Wood R.	0 to 17.9	pH	FallWinterSpring	pH 6.5 to 9.0	Resident fish and aquatic life; Water contact recreation	3
UKL	Wood R.	0 to 17.9	pH	Summer	pH 6.5 to 9.0	Resident fish and aquatic life; Water contact recreation	3
UKL	Wood R.	0 to 17.8	Phosphate Phosphorus	Summer	Total phosphates as phosphorus <sup>4</sup>	Aquatic life	3
UKL	Wood R.	0 to 17.8	Sedimentation	Undefined	Bottom Deposits <sup>2</sup>	Resident fish and aquatic life; Salmonid fish rearing; Salmonid fish spawning	3
UKL	Wood R.	0 to 17.8	Temperature	Summer	Rearing: 17.8 C	Anadromous fish passage; Salmonid fish rearing	Attaining
UKL; Lost	Unnamed	0 to 3.9	Alkalinity	Year Around	Table 20 Toxic Substances	Aquatic life	3
UKL; Lost	Unnamed	0 to 3.9	Ammonia	Year Around	Table 20 Toxic Substances	Aquatic life	3
UKL; Lost	Unnamed	0 to 3.9	Chlorophyll a	Summer	Reservoir, river, estuary, non-thermally stratified lake: 0.015 mg/l	Aesthetics; Fishing; Livestock watering; Water contact recreation; Water supply	3
UKL; Lost	Unnamed	0 to 3.9	Chlorophyll a	FallWinterSpring	Reservoir, river, estuary, non-thermally stratified lake: 0.015 mg/l	Aesthetics; Fishing; Livestock watering; Water contact recreation; Water supply	3
UKL; Lost	Unnamed	0 to 3.9	Dissolved Oxygen	Year Around (Non-spawning)	Cool water: Not less than 6.5 mg/l	Cool-water aquatic life	3
UKL; Lost	Unnamed	0 to 3.9	E. Coli	Summer	E. Coli <sup>3</sup>	Water contact recreation	3
UKL; Lost	Unnamed	0 to 3.9	E. Coli	FallWinterSpring	E. Coli <sup>3</sup>	Water contact recreation	3
UKL; Lost	Unnamed	0 to 3.9	pH	FallWinterSpring	pH 6.5 to 9.0	Resident fish and aquatic life; Water contact recreation	3
UKL; Lost	Unnamed	0 to 3.9	pH	Summer	pH 6.5 to 9.0	Resident fish and aquatic life; Water contact recreation	3
UKL; Lost	Unnamed	0 to 3.9	Phosphate Phosphorus	Summer	Total phosphates as phosphorus <sup>4</sup>	Aquatic life	3
UKL; Lost; Upper Klamath	Klamath R.	207 to 285.3	Alkalinity	Year Around	Table 20 Toxic Substances	Aquatic life	2
UKL; Lost; Upper Klamath	Klamath R.	207 to 285.3	Arsenic	Year Around	Table 40 Human Health Criteria for Toxic Pollutants	Aquatic life; Human health	5
UKL; Lost; Upper Klamath	Klamath R.	207 to 285.3	Chlorophyll a	FallWinterSpring	Reservoir, river, estuary, non-thermally stratified lake: 0.015 mg/l	Aesthetics; Fishing; Livestock watering; Water contact recreation; Water supply	3
UKL; Lost; Upper Klamath	Klamath R.	207 to 285.3	E. Coli	Summer	30-day log mean of 126 E. coli organisms per 100 ml; no single sample > 406 organisms per 100 ml	Water contact recreation	2
UKL; Lost; Upper Klamath	Klamath R.	207 to 285.3	E. Coli	FallWinterSpring	30-day log mean of 126 E. coli organisms per 100 ml; no single	Water contact recreation	2

Watershed (USGS 4th Field Name)	Water Body (Stream/Lake)	River Miles	Parameter	Season	Criteria	Beneficial Uses	Integrated Category
					sample > 406 organisms per 100 ml		
UKL; Lost; Upper Klamath	Klamath R.	207 to 285.3	pH	FallWinterSpring	pH 6.5 to 9.0	Resident fish and aquatic life; Water contact recreation	2
UKL; Lost; Upper Klamath	Klamath R.	207 to 285.3	Phosphate Phosphorus	Summer	Total phosphates as phosphorus <sup>4</sup>	Aquatic life	3B
UKL; Upper Klamath; Lost	Klamath R. / UKL	253 to 275	pH	FallWinterSpring	pH 6.5 to 9.0		5
Williamson	Cottonwood Cr.	0 to 11.9	Temperature	Summer	Rearing: 17.8 C	Anadromous fish passage; Salmonid fish rearing	3B
Williamson	Jackson Cr.	0 to 10.4	Temperature	Summer	Rearing: 17.8 C	Anadromous fish passage; Salmonid fish rearing	Attaining
Williamson	Miller Cr.	0 to 12.7	Biological Criteria	Year Around	Biocriteria <sup>5</sup>	Aquatic life	5
Williamson	Miller Cr.	0 to 12.7	Temperature	Summer	Rearing: 17.8 C	Anadromous fish passage; Salmonid fish rearing	Attaining
Williamson	Sand Cr.	0 to 18	Temperature	Summer	Rearing: 17.8 C	Anadromous fish passage; Salmonid fish rearing	Attaining
Williamson	Tipsoo Cr.	0 to 3	Biological Criteria	Year Around	Biocriteria <sup>5</sup>	Aquatic life	5
Williamson	Williamson R.	12.5 to 94.6	Chlorophyll a	Summer	Reservoir, river, estuary, non- thermally stratified lake: 0.015 mg/l	Aesthetics; Fishing; Livestock watering; Water contact recreation; Water supply	3
Williamson	Williamson R.	12.5 to 94.6	Chlorophyll a	FallWinterSpring	Reservoir, river, estuary, non- thermally stratified lake: 0.015 mg/l	Aesthetics; Fishing; Livestock watering; Water contact recreation; Water supply	3
Williamson	Williamson R.	9.4 to 39.2	Dissolved Oxygen	Year Around (Non-spawning)	Cold water: Not less than 8.0 mg/l or 90% of saturation	Cold-water aquatic life	3
Williamson	Williamson R.	39.2 to 94.6	Dissolved Oxygen	Year Around (Non-spawning)	Cool water: Not less than 6.5 mg/l	Cool-water aquatic life	3
Williamson	Williamson R.	12.5 to 94.6	E. Coli	Summer	E. Coli <sup>3</sup>	Water contact recreation	3
Williamson	Williamson R.	12.5 to 94.6	E. Coli	FallWinterSpring	E. Coli <sup>3</sup>	Water contact recreation	3
Williamson	Williamson R.	35.6 to 94.6	Flow Modification	Undefined	Creation of foul tastes, odors, or toxins. <sup>1</sup>	Resident fish and aquatic life; Salmonid fish rearing; 4B Salmonid fish spawning	
Williamson	Williamson R.	12.5 to 35.6	Flow Modification	Undefined	Creation of foul tastes, odors, or toxins. <sup>1</sup>	Resident fish and aquatic life; Salmonid fish rearing; 4B Salmonid fish spawning	
Williamson	Williamson R.	35.6 to 94.6	Habitat Modification	Undefined	Creation of foul tastes, odors, or toxins. <sup>1</sup>	Resident fish and aquatic life; Salmonid fish rearing; 4B Salmonid fish spawning	
Williamson	Williamson R.	12.5 to 35.6	Habitat Modification	Undefined	Creation of foul tastes, odors, or toxins. <sup>1</sup>	Resident fish and aquatic life; Salmonid fish rearing; 4B Salmonid fish spawning	
Williamson	Williamson R.	35.6 to 94.6	Nutrients	Undefined		Aesthetics	3

<b>Watershed (USGS 4th Field Name)</b>	<b>Water Body (Stream/Lake)</b>	<b>River Miles</b>	<b>Parameter</b>	<b>Season</b>	<b>Criteria</b>	<b>Beneficial Uses</b>	<b>Integrated Category</b>
Williamson	Williamson R.	12.5 to 35.6	Nutrients	Undefined		Aesthetics	3
Williamson	Williamson R.	12.5 to 94.6	pH	FallWinterSpring	pH 6.5 to 9.0	Resident fish and aquatic life; Water contact recreation	3
Williamson	Williamson R.	12.5 to 94.6	pH	Summer	pH 6.5 to 9.0	Resident fish and aquatic life; Water contact recreation	3
Williamson	Williamson R.	0 to 94.6	Sedimentation	Year Around	Bottom Deposits <sup>2</sup>		5
Williamson	Williamson R.	12.5 to 35.6	Sedimentation	Undefined	Bottom Deposits <sup>2</sup>	Resident fish and aquatic life; Salmonid fish rearing; Salmonid fish spawning	3
Williamson	Williamson R.	35.6 to 94.6	Sedimentation	Undefined	Bottom Deposits <sup>2</sup>	Resident fish and aquatic life; Salmonid fish rearing; Salmonid fish spawning	3
Williamson	Williamson R.	35.6 to 94.6	Temperature	Summer	Rearing: 17.8 C	Anadromous fish passage; Salmonid fish rearing	4A
Williamson	Williamson R.	12.5 to 35.6	Temperature	Summer	Rearing: 17.8 C	Anadromous fish passage; Salmonid fish rearing	4A
Williamson; UKL	Williamson R.	0 to 94.6	Alkalinity	Year Around	Table 20 Toxic Substances	Aquatic life	2
Williamson; UKL	Williamson R.	0 to 94.6	Ammonia	Year Around	Table 20 Toxic Substances	Aquatic life	2
Williamson; UKL	Williamson R.	0 to 94.6	Chloride	Year Around	Table 20 Toxic Substances	Aquatic life	3
Williamson; UKL	Williamson R.	0 to 12.5	Chlorophyll a	Summer	Reservoir, river, estuary, non-thermally stratified lake: 0.015 mg/l	Aesthetics; Fishing; Livestock watering; Water contact recreation; Water supply	2
Williamson; UKL	Williamson R.	0 to 12.5	Chlorophyll a	FallWinterSpring	Reservoir, river, estuary, non-thermally stratified lake: 0.015 mg/l	Aesthetics; Fishing; Livestock watering; Water contact recreation; Water supply	3
Williamson; UKL	Williamson R.	0 to 9.4	Dissolved Oxygen	Year Around (Non-spawning)	Cool water: Not less than 6.5 mg/l	Cool-water aquatic life	2
Williamson; UKL	Williamson R.	0 to 94.6	Dissolved Oxygen	January 1 - May 15	Spawning: Not less than 11.0 mg/L or 95% of saturation	Resident trout spawning	5
Williamson; UKL	Williamson R.	0 to 12.5	E. Coli	Summer	30-day log mean of 126 E. coli organisms per 100 ml; no single sample > 406 organisms per 100 ml	Water contact recreation	2
Williamson; UKL	Williamson R.	0 to 12.5	E. Coli	FallWinterSpring	30-day log mean of 126 E. coli organisms per 100 ml; no single sample > 406 organisms per 100 ml	Water contact recreation	2
Williamson; UKL	Williamson R.	0 to 12.5	Fecal Coliform	Summer	Fecal coliform log mean of 200 organisms per 100 ml; no more than 10% > 400 per 100 ml	Water contact recreation	Attaining
Williamson; UKL	Williamson R.	0 to 12.5	Fecal Coliform	FallWinterSpring	Fecal coliform log mean of 200 organisms per 100 ml; no more than 10% > 400 per 100 ml	Water contact recreation	Attaining

<b>Watershed (USGS 4th Field Name)</b>	<b>Water Body (Stream/Lake)</b>	<b>River Miles</b>	<b>Parameter</b>	<b>Season</b>	<b>Criteria</b>	<b>Beneficial Uses</b>	<b>Integrated Category</b>
Williamson; UKL	Williamson R.	0 to 12.5	Flow Modification	Undefined	Creation of foul tastes, odors, or toxins. <sup>1</sup>	Resident fish and aquatic life; Salmonid fish rearing; 4B Salmonid fish spawning	
Williamson; UKL	Williamson R.	0 to 12.5	Habitat Modification	Undefined	Creation of foul tastes, odors, or toxins. <sup>1</sup>	Resident fish and aquatic life; Salmonid fish rearing; 4B Salmonid fish spawning	
Williamson; UKL	Williamson R.	0 to 94.6	Iron	Year Around	Table 20 Toxic Substances	Aquatic life; Human health	3
Williamson; UKL	Williamson R.	0 to 94.6	Manganese	Year Around	Table 20 Toxic Substances	Human health	3
Williamson; UKL	Williamson R.	0 to 12.5	pH	FallWinterSpring	pH 6.5 to 9.0	Anadromous fish passage; Resident fish and aquatic life; Salmonid fish rearing; Salmonid fish spawning; Water contact recreation	2
Williamson; UKL	Williamson R.	0 to 12.5	pH	Summer	pH 6.5 to 9.0	Anadromous fish passage; Resident fish and aquatic life; Salmonid fish rearing; Salmonid fish spawning; Water contact recreation	2
Williamson; UKL	Williamson R.	0 to 94.6	Phosphate Phosphorus	Summer	Total phosphates as phosphorus <sup>4</sup>	Aquatic life	3B
Williamson; UKL	Williamson R.	0 to 12.5	Sedimentation	Undefined	Bottom Deposits <sup>2</sup>	Resident fish and aquatic life; Salmonid fish rearing; 3 Salmonid fish spawning	
Williamson; UKL	Williamson R.	0 to 12.5	Temperature	Summer	Rearing: 17.8 C	Anadromous fish passage; Salmonid fish rearing	4A

<sup>1</sup>The creation of tastes or odors or toxic or other conditions that are deleterious to fish or other aquatic life or affect the potability of drinking water or the palatability of fish or shellfish may not be allowed.

<sup>2</sup>The formation of appreciable bottom or sludge deposits or the formation of any organic or inorganic deposits deleterious to fish or other aquatic life or injurious to public health, recreation, or industry may not be allowed.

<sup>3</sup>30-day log mean of 126 E. coli organisms per 100 ml; no single sample > 406 organisms per 100 ml

<sup>4</sup>Total phosphates as phosphorus (P): Benchmark 50 ug/L in streams to control excessive aquatic growths

<sup>5</sup>Biocriteria: Waters of the state must be of sufficient quality to support aquatic species without detrimental changes in the resident biological communities.

<sup>6</sup>The development of fungi or other growths having a deleterious effect on stream bottoms, fish or other aquatic life, or which are injurious to health, recreation or industry may not be allowed.

Table A11. Oregon Department of Environmental Quality (ODEQ) list of impaired lakes and reservoirs in the Upper Klamath Basin.

Water Body (Stream/Lake)	Parameter	Season	Criteria	Beneficial Uses	Integrated Category
Keene Creek / Hyatt Reservoir	Aquatic Weeds Or Algae	Undefined	Fungi or algae development <sup>1</sup>	Aesthetics; Drinking water; Fishing; Livestock watering; Water contact recreation; Water supply	5
Keene Creek / Hyatt Reservoir	Nutrients	Undefined		Aesthetics	3
Klamath River / Agency Lake	Dissolved Oxygen	Summer	Cool water: Not less than 6.5 mg/l	Resident fish and aquatic life; Salmonid fish rearing	4
Klamath River / Agency Lake	Chlorophyll a	Summer	Thermally stratified lake: 0.01 mg/l	Aesthetics; Fishing; Livestock watering; Water contact recreation; Water supply	4
Klamath River / Agency Lake	Sedimentation	Undefined	Bottom deposits <sup>2</sup>	Resident fish and aquatic life; Salmonid fish rearing; Salmonid fish spawning	3
Klamath River / Agency Lake	Nutrients	Undefined		Aesthetics	3
Klamath River / Agency Lake	Aquatic Weeds Or Algae	Undefined	Fungi or algae development <sup>1</sup>	Aesthetics; Drinking water; Fishing; Livestock watering; Water contact recreation; Water supply	4
Klamath River / Agency Lake	pH	Summer	pH 6.5 to 8.5	Anadromous fish passage; Resident fish and aquatic life; Salmonid fish rearing; Salmonid fish spawning; Water contact recreation	4
Klamath River / Ewauna, Lake	Dissolved Oxygen	Year Around (Non-spawning)	Cool water: Not less than 6.5 mg/l	Cool-water aquatic life	5
Klamath River / Ewauna, Lake	Ammonia	Year Around	Table 20 Toxic Substances		5
Klamath River / Ewauna, Lake	pH	Summer	pH 6.0 to 8.5		5
Klamath River / Ewauna, Lake	pH	FallWinterSpring	pH 6.0 to 8.5		5
Klamath River / Ewauna, Lake	Phosphate Phosphorus	Summer	Total phosphates as phosphorus (P): Benchmark 50 ug/L in streams to control excessive aquatic growths	Aquatic life	3B
Klamath River / J.C. Boyle Reservoir	pH	Summer	pH 6.0 to 8.5	Resident fish and aquatic life; Water contact recreation	3
Klamath River / J.C. Boyle Reservoir	Chlorophyll a	Summer	Reservoir, river, estuary, non-thermally stratified lake: 0.015 mg/l	Fishing; Aesthetics; Livestock watering; Water contact recreation; Water supply	3
Klamath River / J.C. Boyle Reservoir	Nutrients	Undefined		Aesthetics	3

<b>Water Body (Stream/Lake)</b>	<b>Parameter</b>	<b>Season</b>	<b>Criteria</b>	<b>Beneficial Uses</b>	<b>Integrated Category</b>
Klamath River / Upper Klamath Lake	Chlorophyll a	Summer	Thermally stratified lake: 0.01 mg/l	Aesthetics; Fishing; Livestock watering; Water contact recreation; Water supply	4
Klamath River / Upper Klamath Lake	Sedimentation	Undefined	Bottom deposits <sup>2</sup>	Resident fish and aquatic life; Salmonid fish rearing; Salmonid fish spawning	3
Klamath River / Upper Klamath Lake	pH	FallWinterSpring	pH 6.5 to 9.0		5
Klamath River / Upper Klamath Lake	Nutrients	Undefined		Aesthetics	3
Klamath River / Upper Klamath Lake	pH	Summer	pH 6.5 to 8.5	Anadromous fish passage; Resident fish and aquatic life; Salmonid fish rearing; Salmonid fish spawning; Water contact recreation	4
Klamath River / Upper Klamath Lake	Dissolved Oxygen	Summer	Cool water: Not less than 6.5 mg/l	Resident fish and aquatic life; Salmonid fish rearing	4
Klamath River / Upper Klamath Lake	Aquatic Weeds Or Algae	Undefined	Fungi or algae development <sup>1</sup>	Aesthetics; Drinking water; Fishing; Livestock watering; Water contact recreation; Water supply	4
Klamath River / Upper Klamath Lake	Temperature	Undefined		Anadromous fish passage; Resident fish and aquatic life; Salmonid fish rearing; Salmonid fish spawning	3
Obenchain Reservoir	Aquatic Weeds Or Algae	Undefined	Fungi or algae development <sup>1</sup>	Aesthetics; Fishing; Water contact recreation	3
Obenchain Reservoir	Dissolved Oxygen	Undefined		Anadromous fish passage; Resident fish and aquatic life; Salmonid fish rearing; Salmonid fish spawning	3
Obenchain Reservoir	Nutrients	Undefined		Aesthetics	3

<sup>1</sup>The development of fungi or other growths having a deleterious effect on stream bottoms, fish or other aquatic life, or which are injurious to health, recreation or industry may not be allowed.

<sup>2</sup>The formation of appreciable bottom or sludge deposits or the formation of any organic or inorganic deposits deleterious to fish or other aquatic life or injurious to public health, recreation, or industry may not be allowed.

## APPENDIX B: CALIFORNIA LIST OF IMPAIRED WATERBODIES

Table B12. California 303(d) list for the portion of the Klamath River from the Oregon border downstream to Scott River, USEPA approved 2012. Table modified from California Integrated Report online combined 303(d) list [http://www.waterboards.ca.gov/water\\_issues/programs/tmdl/integrated2012.shtml](http://www.waterboards.ca.gov/water_issues/programs/tmdl/integrated2012.shtml).

WATER BODY NAME	POLLUTANT	POLLUTANT CATEGORY	INTEGRATED REPORT CATEGORY	POTENTIAL SOURCES	SOURCE CATEGORY
Klamath River HU, Middle HA, Iron Gate Dam to Scott River <sup>1</sup>	Aluminum	Metals/Metalloids	5	Source Unknown	Source Unknown
Klamath River HU, Middle HA, Iron Gate Dam to Scott River <sup>2</sup>	Cyanobacteria hepatotoxic microcystins	Miscellaneous	5	Agriculture	Agriculture
Klamath River HU, Middle HA, Iron Gate Dam to Scott River <sup>2</sup>	Cyanobacteria hepatotoxic microcystins	Miscellaneous	5	Grazing-Related Sources	Agriculture
Klamath River HU, Middle HA, Iron Gate Dam to Scott River <sup>2</sup>	Cyanobacteria hepatotoxic microcystins	Miscellaneous	5	Irrigated Crop Production	Agriculture
Klamath River HU, Middle HA, Iron Gate Dam to Scott River	Nutrients	Nutrients	5	Agriculture	Agriculture
Klamath River HU, Middle HA, Iron Gate Dam to Scott River	Nutrients	Nutrients	5	Grazing-Related Sources	Agriculture
Klamath River HU, Middle HA, Iron Gate Dam to Scott River	Nutrients	Nutrients	5	Irrigated Crop Production	Agriculture
Klamath River HU, Middle HA, Iron Gate Dam to Scott River <sup>1</sup>	Organic Enrichment/Low Dissolved Oxygen	Nutrients	5	Agriculture	Agriculture
Klamath River HU, Middle HA, Iron Gate Dam to Scott River <sup>1</sup>	Organic Enrichment/Low Dissolved Oxygen	Nutrients	5	Grazing-Related Sources	Agriculture
Klamath River HU, Middle HA, Iron Gate Dam to Scott River <sup>1</sup>	Organic Enrichment/Low Dissolved Oxygen	Nutrients	5	Irrigated Crop Production	Agriculture
Klamath River HU, Middle HA, Iron Gate Dam to Scott River	Temperature, water	Miscellaneous	5	Agriculture	Agriculture
Klamath River HU, Middle HA, Iron Gate Dam to Scott River	Temperature, water	Miscellaneous	5	Grazing-Related Sources	Agriculture
Klamath River HU, Middle HA, Iron Gate Dam to Scott River	Temperature, water	Miscellaneous	5	Irrigated Crop Production	Agriculture
Klamath River HU, Middle HA, Iron Gate Dam to Scott River <sup>2</sup>	Cyanobacteria hepatotoxic microcystins	Miscellaneous	5	Road Construction	Construction/Land Development
Klamath River HU, Middle HA, Iron Gate Dam to Scott River	Nutrients	Nutrients	5	Road Construction	Construction/Land Development
Klamath River HU, Middle HA, Iron Gate Dam to Scott River <sup>1</sup>	Organic Enrichment/Low Dissolved Oxygen	Nutrients	5	Road Construction	Construction/Land Development
Klamath River HU, Middle HA, Iron Gate Dam to Scott River	Temperature, water	Miscellaneous	5	Road Construction	Construction/Land Development
Klamath River HU, Middle HA, Iron Gate Dam to Scott River <sup>2</sup>	Cyanobacteria hepatotoxic microcystins	Miscellaneous	5	Habitat Modification	Habitat Modification
Klamath River HU, Middle HA, Iron Gate Dam to Scott River	Nutrients	Nutrients	5	Drainage/Filling Of Wetlands	Habitat Modification
Klamath River HU, Middle HA, Iron Gate Dam to Scott River <sup>1</sup>	Organic Enrichment/Low Dissolved Oxygen	Nutrients	5	Drainage/Filling Of Wetlands	Habitat Modification
Klamath River HU, Middle HA, Iron Gate Dam to Scott River	Temperature, water	Miscellaneous	5	Drainage/Filling Of Wetlands	Habitat Modification
Klamath River HU, Middle HA, Iron Gate Dam to Scott River	Temperature, water	Miscellaneous	5	Removal of Riparian Vegetation	Habitat Modification
Klamath River HU, Middle HA, Iron Gate Dam to Scott River	Temperature, water	Miscellaneous	5	Flow Alteration/Regulation/Modification	Hydromodification

WATER BODY NAME	POLLUTANT	POLLUTANT CATEGORY	INTEGRATED REPORT CATEGORY	POTENTIAL SOURCES	SOURCE CATEGORY
Klamath River HU, Middle HA, Iron Gate Dam to Scott River <sup>2</sup>	Cyanobacteria hepatotoxic microcystins	Miscellaneous	5	Logging Road Construction/Maintenance	Silviculture
Klamath River HU, Middle HA, Iron Gate Dam to Scott River <sup>2</sup>	Cyanobacteria hepatotoxic microcystins	Miscellaneous	5	Silviculture	Silviculture
Klamath River HU, Middle HA, Iron Gate Dam to Scott River	Nutrients	Nutrients	5	Logging Road Construction/Maintenance	Silviculture
Klamath River HU, Middle HA, Iron Gate Dam to Scott River	Nutrients	Nutrients	5	Silviculture	Silviculture
Klamath River HU, Middle HA, Iron Gate Dam to Scott River <sup>1</sup>	Organic Enrichment/Low Dissolved Oxygen	Nutrients	5	Logging Road Construction/Maintenance	Silviculture
Klamath River HU, Middle HA, Iron Gate Dam to Scott River <sup>1</sup>	Organic Enrichment/Low Dissolved Oxygen	Nutrients	5	Silviculture	Silviculture
Klamath River HU, Middle HA, Iron Gate Dam to Scott River	Temperature, water	Miscellaneous	5	Logging Road Construction/Maintenance	Silviculture
Klamath River HU, Middle HA, Iron Gate Dam to Scott River	Temperature, water	Miscellaneous	5	Silviculture	Silviculture
Klamath River HU, Middle HA, Oregon to Iron Gate <sup>3</sup>	Temperature, water	Miscellaneous	4a	Other	Source Unknown
Klamath River HU, Middle HA, Oregon to Iron Gate <sup>4</sup>	Cyanobacteria hepatotoxic microcystins	Miscellaneous	4a	Agriculture	Agriculture
Klamath River HU, Middle HA, Oregon to Iron Gate <sup>4</sup>	Cyanobacteria hepatotoxic microcystins	Miscellaneous	4a	Grazing-Related Sources	Agriculture
Klamath River HU, Middle HA, Oregon to Iron Gate <sup>4</sup>	Cyanobacteria hepatotoxic microcystins	Miscellaneous	4a	Irrigated Crop Production	Agriculture
Klamath River HU, Middle HA, Oregon to Iron Gate	Nutrients	Nutrients	4a	Agriculture	Agriculture
Klamath River HU, Middle HA, Oregon to Iron Gate	Nutrients	Nutrients	4a	Grazing-Related Sources	Agriculture
Klamath River HU, Middle HA, Oregon to Iron Gate	Nutrients	Nutrients	4a	Irrigated Crop Production	Agriculture
Klamath River HU, Middle HA, Oregon to Iron Gate <sup>1</sup>	Organic Enrichment/Low Dissolved Oxygen	Nutrients	4a	Agriculture	Agriculture
Klamath River HU, Middle HA, Oregon to Iron Gate <sup>1</sup>	Organic Enrichment/Low Dissolved Oxygen	Nutrients	4a	Grazing-Related Sources	Agriculture
Klamath River HU, Middle HA, Oregon to Iron Gate <sup>1</sup>	Organic Enrichment/Low Dissolved Oxygen	Nutrients	4a	Irrigated Crop Production	Agriculture
Klamath River HU, Middle HA, Oregon to Iron Gate <sup>3</sup>	Temperature, water	Miscellaneous	4a	Grazing-Related Sources	Agriculture
Klamath River HU, Middle HA, Oregon to Iron Gate <sup>3</sup>	Temperature, water	Miscellaneous	4a	Irrigated Crop Production	Agriculture
Klamath River HU, Middle HA, Oregon to Iron Gate <sup>4</sup>	Cyanobacteria hepatotoxic microcystins	Miscellaneous	4a	Road Construction	Construction/Land Development
Klamath River HU, Middle HA, Oregon to Iron Gate	Nutrients	Nutrients	4a	Road Construction	Construction/Land Development
Klamath River HU, Middle HA, Oregon to Iron Gate <sup>1</sup>	Organic Enrichment/Low Dissolved Oxygen	Nutrients	4a	Road Construction	Construction/Land Development
Klamath River HU, Middle HA, Oregon to Iron Gate <sup>3</sup>	Temperature, water	Miscellaneous	4a	Road Construction	Construction/Land Development
Klamath River HU, Middle HA, Oregon to Iron Gate <sup>4</sup>	Cyanobacteria hepatotoxic microcystins	Miscellaneous	4a	Drainage/Filling Of Wetlands	Habitat Modification



WATER BODY NAME	POLLUTANT	POLLUTANT CATEGORY	INTEGRATED REPORT CATEGORY	POTENTIAL SOURCES	SOURCE CATEGORY
Klamath River HU, Middle HA, Oregon to Iron Gate	Nutrients	Nutrients	4a	Drainage/Filling Of Wetlands	Habitat Modification
Klamath River HU, Middle HA, Oregon to Iron Gate <sup>1</sup>	Organic Enrichment/Low Dissolved Oxygen	Nutrients	4a	Drainage/Filling Of Wetlands	Habitat Modification
Klamath River HU, Middle HA, Oregon to Iron Gate <sup>3</sup>	Temperature, water	Miscellaneous	4a	Removal of Riparian Vegetation	Habitat Modification
Klamath River HU, Middle HA, Oregon to Iron Gate <sup>3</sup>	Temperature, water	Miscellaneous	4a	Flow Alteration/Regulation/Modification	Hydromodification
Klamath River HU, Middle HA, Oregon to Iron Gate <sup>3</sup>	Temperature, water	Miscellaneous	4a	Sediment Resuspension (clean sediment)	Sediment
Klamath River HU, Middle HA, Oregon to Iron Gate <sup>4</sup>	Cyanobacteria hepatotoxic microcystins	Miscellaneous	4a	Logging Road Construction/Maintenance	Silviculture
Klamath River HU, Middle HA, Oregon to Iron Gate <sup>4</sup>	Cyanobacteria hepatotoxic microcystins	Miscellaneous	4a	Silviculture	Silviculture
Klamath River HU, Middle HA, Oregon to Iron Gate	Nutrients	Nutrients	4a	Logging Road Construction/Maintenance	Silviculture
Klamath River HU, Middle HA, Oregon to Iron Gate	Nutrients	Nutrients	4a	Silviculture	Silviculture
Klamath River HU, Middle HA, Oregon to Iron Gate <sup>1</sup>	Organic Enrichment/Low Dissolved Oxygen	Nutrients	4a	Logging Road Construction/Maintenance	Silviculture
Klamath River HU, Middle HA, Oregon to Iron Gate <sup>1</sup>	Organic Enrichment/Low Dissolved Oxygen	Nutrients	4a	Silviculture	Silviculture
Klamath River HU, Middle HA, Oregon to Iron Gate <sup>3</sup>	Temperature, water	Miscellaneous	4a	Logging Road Construction/Maintenance	Silviculture
Klamath River HU, Middle HA, Oregon to Iron Gate <sup>3</sup>	Temperature, water	Miscellaneous	4a	Silviculture	Silviculture
Klamath River HU, Middle HA, Oregon to Iron Gate	Nutrients	Nutrients	4a	Nonpoint Source	Unspecified Nonpoint Source

<sup>1</sup>This listing only applies to the portion of the mainstem Klamath River that lies within the Klamath River HU, Middle HA, Iron Gate Dam to Scott River water body.

<sup>2</sup>This listing applies to the mainstem Klamath River in the Klamath River Hydrologic Unit, Middle Klamath River Hydrologic Area, Iron Gate Dam to Scott River reach.

<sup>3</sup>The Klamath River HU, Middle HA, Oregon to Iron Gate Dam includes the following Hydrologic Sub Areas (HSAs): Iron Gate HSA 115.37 and Copco HSA 105.38.

<sup>4</sup>This listing applies to the mainstem Klamath River in the Klamath River Hydrologic Unit, Middle Klamath River Hydrologic Area, Oregon to Iron Gate reach, excluding the riverine reach from the Oregon border downstream to the beginning of Copco 1 Reservoir.

<b>WATER BODY NAME</b>	<b>POLLUTANT</b>	<b>POLLUTANT CATEGORY</b>	<b>INTEGRATED REPORT CATEGORY</b>	<b>POTENTIAL SOURCES</b>	<b>SOURCE CATEGORY</b>
Copco Lake <sup>1</sup>	Mercury	Metals/Metalloids	5	Source Unknown	Source Unknown
Copco Lake <sup>2</sup>	Cyanobacteria hepatotoxic microcystins	Miscellaneous	5	Agriculture	Agriculture
Copco Lake <sup>2</sup>	Cyanobacteria hepatotoxic microcystins	Miscellaneous	5	Grazing-Related Sources	Agriculture
Copco Lake <sup>2</sup>	Cyanobacteria hepatotoxic microcystins	Miscellaneous	5	Road Construction	Construction/Land Development
Copco Lake <sup>2</sup>	Cyanobacteria hepatotoxic microcystins	Miscellaneous	5	Drainage/Filling Of Wetlands	Habitat Modification
Copco Lake <sup>2</sup>	Cyanobacteria hepatotoxic microcystins	Miscellaneous	5	Silviculture	Silviculture
Iron Gate Reservoir	Mercury	Metals/Metalloids	5	Source Unknown	Source Unknown
Iron Gate Reservoir	Cyanobacteria hepatotoxic microcystins	Miscellaneous	5	Agriculture	Agriculture
Iron Gate Reservoir	Cyanobacteria hepatotoxic microcystins	Miscellaneous	5	Grazing-Related Sources	Agriculture
Iron Gate Reservoir	Cyanobacteria hepatotoxic microcystins	Miscellaneous	5	Irrigated Crop Production	Agriculture
Iron Gate Reservoir	Cyanobacteria hepatotoxic microcystins	Miscellaneous	5	Road Construction	Construction/Land Development
Iron Gate Reservoir	Cyanobacteria hepatotoxic microcystins	Miscellaneous	5	Drainage/Filling Of Wetlands	Habitat Modification
Iron Gate Reservoir	Cyanobacteria hepatotoxic microcystins	Miscellaneous	5	Logging Road Construction/Maintenance	Silviculture
Iron Gate Reservoir	Cyanobacteria hepatotoxic microcystins	Miscellaneous	5	Silviculture	Silviculture
Klamath River HU, Tule Lake and Lower Klamath Lake National Wildlife Refuge	pH (high)	Miscellaneous	4a	Internal Nutrient Cycling (primarily lakes)	Natural Sources
Klamath River HU, Tule Lake and Lower Klamath Lake National Wildlife Refuge	pH (high)	Miscellaneous	4a	Nonpoint Source	Unspecified Nonpoint Source

<sup>1</sup>This listing applies to Copco 1 Reservoir.

<sup>2</sup>This listing applies to the Copco 1 and Copco 2 Reservoirs.